

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

EP 0 874 045 A1

(12)

**EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

(43) Date of publication:

28.10.1998 Bulletin 1998/44

(21) Application number: 97935810.8

(22) Date of filing: 19.08.1997

(51) Int. Cl.<sup>6</sup>: C12N 15/00, C12P 21/00

(86) International application number:

PCT/JP97/02859

(87) International publication number:

WO 98/07840 (26.02.1998 Gazette 1998/08)

(84) Designated Contracting States:

AT BE CH DE DK ES FR GB IE IT LI LU NL SE

(30) Priority: 19.08.1996 JP 235928/96

(83) Declaration under Rule 28(4) EPC (expert solution)

(71) Applicant:

SNOW BRAND MILK PRODUCTS CO., LTD.  
Sapporo-shi, Hokkaido 065 (JP)

(72) Inventors:

- NAKAGAWA, Nobuaki,  
Nishiura Heights 2-4  
Shimotsuga-gun, Tochigi 329-05 (JP)
- YASUDA, Hisataka  
Kawachi-gun, Tochigi 329-04 (JP)
- MORINAGA, Tomonori  
Shimotsuga-gun, Tochigi 321-02 (JP)

(74) Representative:

Wakerley, Helen Rachael  
Reddie & Grose,  
16 Theobalds Road  
London WC1X 8PL (GB)(54) **NOVEL DNAs AND PROCESS FOR PRODUCING PROTEINS BY USING THE SAME**

(57) DNAs having the nucleotide sequences of the Sequences No. 1 and No. 2 in the Sequence Table and a process for producing a protein which comprises inserting these DNAs into expression vectors to thereby produce a protein having molecular weights of about 60 kD (under reductive conditions) and about 60 kD and 120 kD (under non-reductive conditions) and being capable of inhibiting formation of osteoclast. These proteins are useful in the treatment of osteoporosis and rheumatism.

EP 0 874 045 A1

## Description

FIELD OF TECHNOLOGY

5 The present invention relates to a novel DNA and a process for preparing a protein which possesses an activity to inhibit osteoclast differentiation and/or maturation (hereinafter called osteoclastogenesis-inhibitory activity) by a genetic engineering technique using the DNA. More particularly, the present invention relates to a genomic DNA encoding a protein OCIF which possesses an osteoclastogenesis-inhibitory activity and a process for preparing said protein by a genetic engineering technique using the genomic DNA.

BACKGROUND OF THE INVENTION

Human bones are constantly repeating a process of resorption and formation. Osteoclasts controlling formation of bones and osteoclasts controlling resorption of bones take major roles in this process. Osteoporosis is a typical disease caused by abnormal metabolism of bones. This disease is caused when bone resorption by osteoclasts exceeds bone formation by osteoblasts. Although the mechanism of this disease is still to be elucidated completely, the disease causes the bones to ache, makes the bones fragile, and may result in fracturing of the bones. As the population of the aged increases, this disease results in an increase in bedridden aged people which becomes a social problem. Urgent development of a therapeutic agent for this disease is strongly desired. Disease due to a decrease in bone mass is expected to be treated by controlling bone resorption, accelerating bone formation, or improving balance between bone resorption and formation.

Osteogenesis is expected to increase by accelerating proliferation, differentiation, or activation of the cells controlling bone formation, or by controlling proliferation, differentiation, or activation of the cells involved in bone resorption. In recent years, strong interest has been directed to physiologically active proteins (cytokines) exhibiting such activities as described above, and energetic research is ongoing on this subject. The cytokines which have been reported to accelerate proliferation or differentiation of osteoblasts include the proteins of fibroblast growth factor family (FGF: Rodan S. B. et al., *Endocrinology* vol. 121, p1917, 1987), insulin-like growth factor I (IGF-I: Hock J. M. et al., *Endocrinology* vol. 122, p 254, 1988), insulin growth factor II (IGF-II: McCarthy T. et al., *Endocrinology* vol. 124, p 301, 1989), Activin A (Centrella M. et al., *Mol. Cell. Biol.*, vol. 11, p 250, 1991), transforming growth factor- $\beta$  (Noda M., *The Bone*, vol. 2, p 29, 1988), Vascutotropin (Varonique M. et al., *Biochem. Biophys. Res. Commun.*, vol. 199, p 380, 1994), and the protein of heterotopic bone formation factor family (bone morphogenic protein; BMP: BMP-2: Yanaguchi A. et al., *J. Cell Biol.* vol. 113, p 682, 1991, OP-1: Sampath T. K. et al., *J. Biol. Chem.* vol. 267, p 20532, 1992, and Knutsen R. et al., *Biochem. Biophys. Res. Commun.* vol. 194, P 1352, 1993).

On the other hand, as the cytokines which suppress differentiation and/or maturation of osteoclasts, transforming growth factor- $\beta$  (Chen C. et al., *Proc. Natl. Acad. Sci. USA*, vol. 85, p 5683, 1988), interleukin-4 (Kasano K. et al., *Bone-Miner.*, vol. 21, p 179, 1993), and the like have been reported. Further, as the cytokines which suppress bone resorption by osteoclast, calcitonin (*Bone-Miner.*, vol. 17, p 347, 1992), macrophage colony stimulating factor (Hattersley G. et al., *J. Cell. Physiol.* vol. 137, p 199, 1988), interleukin-4 (Watanabe, K. et al., *Biochem. Biophys. Res. Commun.* vol. 172, P 1035, 1990), and interferon- $\gamma$  (Gowen M. et al., *J. Bone Miner. Res.*, vol. 1, p 46.9, 1986) have been reported.

These cytokines are expected to be used as agents for treating diseases accompanying bone loss by accelerating bone formation or suppressing of bone resorption. Clinical tests are being undertaken to verify the effect of improving bone metabolism of some cytokines such as insulin-like growth factor-I and the heterotopic bone formation factor family. In addition, calcitonin is already commercially available as a therapeutic agent for osteoporosis and a pain relief agent. At present, drugs for clinically treating bone diseases or shortening the period of treatment of bone diseases include activated vitamin D<sub>3</sub>, calcitonin and its derivatives, and hormone preparations such as estradiol agent, ipriflavon or calcium preparations. These agents are not necessarily satisfactory in terms of the efficacy and therapeutic results. Development of a novel therapeutic agent which can be used in place of these agents is strongly desired.

In view of this situation, the present inventors have undertaken extensive studies. As a result, the present inventors had found protein OCIF exhibiting an osteoclastogenesis-inhibitory activity in a culture broth of human embryonic lung fibroblast IMR-90 (ATCC Deposition No. CCL186), and filed a patent application (PCT/JP96/00374). The present inventors have conducted further studies relating to the origin of this protein OCIF exhibiting the osteoclastogenesis-inhibitory activity. The studies have matured into determination of the sequence of a genomic DNA encoding the human origin OCIF. Accordingly, an object of the present invention is to provide a genomic DNA encoding protein OCIF exhibiting osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA.

DISCLOSURE OF THE INVENTION

Specifically, the present invention relates to a genomic DNA encoding protein OCIF exhibiting osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA. The DNA of the present invention includes the nucleotide sequences No. 1 and No. 2 in the Sequence Table attached hereto.

Moreover, the present invention relates to a process for preparing a protein, comprising inserting a DNA including the nucleotide sequences of the sequences No. 1 and No. 2 in the Sequence Table into an expression vector, producing a vector capable of expressing a protein having the following physicochemical characteristics and exhibiting the activity of inhibiting differentiation and/or maturation of osteoclasts, and producing this protein by a genetic engineering technique,

(a) molecular weight (SDS-PAGE):

- (i) Under reducing conditions: about 60 kD,
- (ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 of the Sequence Table,

(c) affinity:

exhibits affinity to a cation exchanger and heparin, and

(d) thermal stability:

- (i) the osteoclast differentiation and/or maturation inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,
- (ii) the osteoclast differentiation and/or maturation inhibitory activity is lost when treated with heat at 90°C for 10 minutes.

The protein obtained by expressing the gene of the present invention exhibits an osteoclastogenesis-inhibitory activity. This protein is effective as an agent for the treatment and improvement of diseases involving decrease in the amount of bone such as osteoporosis, diseases relating to bone metabolism abnormality such as rheumatism, degenerative joint disease, or multiple myeloma, and is useful as an antigen to establish an immunological diagnosis of such diseases.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a result of Western Blotting analysis of the protein obtained by causing genomic DNA of the present invention to express a protein in Example 4 (ii), wherein lane 1 indicates a marker, lane 2 indicates the culture broth of COS7 cells in which a vector pWESRaOCIF (Example 4 (iii)) has been transfected, and lane 3 is the culture broth of COS7 cell in which a vector pWESRa(control) has been transfected.

BEST MODE FOR CARRYING OUT THE INVENTION

The genomic DNA encoding the protein OCIF which exhibits osteoclastogenesis-inhibitory activity in the present invention can be obtained by preparing a cosmid library using a human placenta genomic DNA and a cosmid vector and by screening this library using DNA fragments which are prepared based on the OCIF cDNA as a probe. The thus-obtained genomic DNA is inserted into a suitable expression vector to prepare an OCIF expression cosmid. A recombinant type OCIF can be obtained by transfecting the genomic DNA into a host organism such as various types of cells or microorganism strains and causing the DNA to express a protein by a conventional method. The resultant protein exhibiting osteoclastogenesis-inhibitory activity (an osteoclastogenesis-inhibitory factor) is useful as an agent for the treatment and improvement of diseases involving a decrease in bone mass such as osteoporosis and other diseases relating to bone metabolism abnormality and also as an antigen to prepare antibodies for establishing immunological diagnosis of such diseases. The protein of the present invention can be prepared as a drug composition for oral or non-oral administration. Specifically, the drug composition of the present invention containing the protein which is an osteoclastogenesis-inhibitory factor as an active ingredient can be safely administered to humans and animals. As the form of drug composition, a composition for injection, composition for intravenous drip, suppository, nasal agent, sublingual agent, percutaneous absorption agent, and the like are given. In the case of the composition for injection, such a composition is a mixture of a pharmacologically effective amount of osteoclastogenesis-inhibitory factor of the present

invention and a pharmaceutically acceptable carrier. The composition may further comprise amino acids, saccharides, cellulose derivatives, and other excipients and/or activation agents, including other organic compounds and inorganic compounds which are commonly added to a composition for injection. When an injection preparation is prepared using the osteoclastogenesis-inhibitory factor of the present invention and these excipients and activation agents, a pH adjuster, buffering agent, stabilizer, solubilizing agent, and the like may be added if necessary to prepare various types of injection agents.

The present invention will now be described in more detail by way of examples which are given for the purpose of illustration and not intended to be limiting of the present invention.

#### Example 1

##### (Preparation of a cosmid library)

A cosmid library was prepared using human placenta genomic DNA (Clontech; Cat. No. 6550-2) and pWE15 cosmid vector (Stratagene). The experiment was carried out following principally the protocol attached to the pWE15 cosmid vector kit of Stratagene Company, provided Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Laboratory (1989)) was referred to for common procedures for handling DNA, E. coli, and phage.

##### (i) Preparation of restrictive enzymolysate of human-genomic DNA

Human placenta genomic DNA dissolved in 750  $\mu$ l of a solution containing 10 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, and 100 mM NaCl was added to four 1.5 ml Eppendorf tubes (tube A, B, C, and D) in the amount of 100  $\mu$ g each. Restriction enzyme MboI was added to these tubes in the amounts of 0.2 unit for tube A, 0.4 unit for tube B, 0.6 unit for tube C, and 0.8 unit for tube D, and DNA was digested for 1 hour. Then, EDTA in the amount to make a 20 mM concentration was added to each tube to terminate the reaction, followed by extraction with phenol/chloroform (1:1). A two-fold amount of ethanol was added to the aqueous layer to precipitate DNA. DNA was collected by centrifugation, washed with 70% ethanol, and DNA in each tube was dissolved in 100  $\mu$ l of TE (10 mM HCl (pH 8.0) + 1 mM EDTA buffer solution, hereinafter called TE). DNA in four tubes was combined in one tube and incubated for 10 minutes at 68°C. After cooling to room temperature, the mixture was overlaid onto a 10%-40 % linear sucrose gradient which was prepared in a buffer containing 20 mM Tris-HCl (pH 8.0), 5 mM EDTA, and 1 mM NaCl in a centrifugal tube (38 ml). The tube was centrifuged at 26,000 rpm for 24 hours at 20°C using a rotor SRP28SA manufactured by Hitachi, Ltd. and 0.4 ml fractions of the sucrose gradient was collected using a fraction collector. A portion of each fraction was subjected to 0.4% agarose electrophoresis to confirm the size of DNA. Fractions containing DNA with a length of 30 kb (kilo base pair) to 40 kb were thus combined. The DNA solution was diluted with TE to make a sucrose concentration to 10% or less and 2.5-fold volumes of ethanol was added to precipitate DNA. DNA was dissolved in TE and stored at 4°C.

##### (ii) Preparation of cosmid vector

The pWE15 cosmid vector obtained from Stratagene Company was completely digested with restriction enzyme BamHI according to the protocol attached to the cosmid vector kit. DNA collected by ethanol precipitation was dissolved in TE to a concentration of 1 mg/ml. Phosphoric acid at the 5'-end of this DNA was removed using calf small intestine alkaline phosphatase, and DNA was collected by phenol extraction and ethanol precipitation. The DNA was dissolved in TE to a concentration of 1 mg/ml.

##### (iii) Ligation of genomic DNA to vector and in vitro packaging

1.5 micrograms of genomic DNA fractionated according to size and 3  $\mu$ g of pWE15 cosmid vector which was digested with restriction enzyme BamHI were ligated in 20  $\mu$ l of a reaction solution using Ready-To-Go T4DNA ligase of Pharmacia Company. The ligated DNA was packaged in vitro using Gigapack™ II packaging extract (Stratagene) according to the protocol. After the packaging reaction, a portion of the reaction mixture was diluted stepwise with an SM buffer solution and mixed with E. coli XL1-Blue MR (Stratagene) which was suspended in 10 mM MgCl<sub>2</sub> to cause phage to infect, and plated onto LB agar plates containing 50  $\mu$ g/ml of ampicillin. The number of colonies produced was counted. The number of colonies per 1  $\mu$ l of packaging reaction was calculated based on this result.

##### (iv) Preparation of a cosmid library

The packaging reaction solution thus prepared was mixed with E. coli XL1-Blue MR and the mixture was plated onto agarose plates containing ampicillin so as to produce 50,000 colonies per agarose plate having a 15 cm of diam-

eter. After incubating the plate overnight at 37°C, an LB culture medium was added in the amount of 3 ml per plate to suspend and collect colonies of *E. coli*. Each agarose plate was again washed with 3 ml of the LB culture medium and the washing was combined with the original suspension of *E. coli*. The *E. coli* collected from all agarose plates was placed in a centrifugal tube, glycerol was added to a concentration of 20%, and ampicillin was further added to make a final concentration of 50 µg/ml. A portion of the *E. coli* suspension was removed and the remainder was stored at -80°C. The removed *E. coli* was diluted stepwise and plated onto an agar plates to count the number of colonies per 1 ml of suspension.

#### Example 2

##### (Screening of cosmid library and purification of colony)

A nitrocellulose filter (Millipore) with a diameter of 14.2 cm was placed on each LB agarose plate with a diameter of 15 cm which contained 50 µg/ml of ampicillin. The cosmid library was plated onto the plates so as to produce 50,000 colonies of *E. coli* per plate, followed by incubation overnight at 37°C. *E. coli* on the nitrocellulose filter was transferred to another nitrocellulose filter according to a conventional method to obtain two replica filters. According to the protocol attached to the cosmid vector kit, cosmid DNA in the *E. coli* on the replica filters was denatured with an alkali, neutralized, and immobilized on the nitrocellulose filter using a Stratalinker (Stratagene). The filters were heated for two hours at 80°C in a vacuum oven. The nitrocellulose filters thus obtained were hybridized using two kinds of DNA produced, respectively, from 5'-end and 3'-end of human OCIF cDNA as probes. Namely, a plasmid was purified from *E. coli* pKB/OCIF10 (deposited at The Ministry of International Trade and Industry, the Agency of Industrial Science and Technology, Biotechnology Laboratory, Deposition No. FERM BP-5267) containing OCIF cDNA. The plasmid containing OCIF cDNA was digested with restriction enzymes KpnI and EcoRI. Fragments thus obtained was separated using agarose gel electrophoresis. KpnI/EcoRI fragment with a length of 0.2 kb was purified using a QIAEX II gel extraction kit (Qiagen). This DNA was labeled with <sup>32</sup>P using the Megaprime DNA Labeling System (Amasham) (5'-DNA probe). Apart from this, a BamHI/EcoRV fragment with a length of 0.2 kb which was produced from the above plasmid by digestion with restriction enzymes BamHI and EcoRV was purified and labeled with <sup>32</sup>P (3'-DNA probe). One of the replica filters described above was hybridized with the 5'-DNA probe and the other with the 3'-DNA probe. Hybridization and washing of the filters were carried out according to the protocol attached to the cosmid vector kit. Autoradiography detected several positive signals with each probe. One colony which gave positive signals with both probe was identified. The colony on the agar plate, which corresponding to the signal on the autoradiogram was isolated and purified. A cosmid was prepared from the purified colony by a conventional method. This cosmid was named pWEOCIF. The size of human genomic DNA contained in this cosmid was about 38 kb.

#### Example 3

##### (Determination of the nucleotide sequence of human OCIF genomic DNA)

##### (i) Subcloning of OCIF genomic DNA

Cosmid pWEOCIF was digested with restriction enzyme EcoRI. After the separation of the DNA fragments thus produced by electrophoresis using a 0.7% agarose gel, the DNA fragments were transferred to a nylon membrane (Hybond-N, Amasham) by the Southern blot technique and immobilized on the nylon membrane using Stratalinker (Stratagene). On the other hand, plasmid pBKOCIF was digested with restriction enzyme EcoRI and a 1.6 kb fragment containing human OCIF cDNA was isolated by agarose gel electrophoresis. The fragment was labeled with <sup>32</sup>P using the Megaprime DNA labeling system (Amasham).

Hybridization of the nylon membranes described above with the <sup>32</sup>P-labeled 1.6-kb OCIF cDNA was performed according to a conventional method detected that DNA fragments with a size of 6 kb, 4 kb, 3.6 kb, and 2.6 kb. These fragments hybridized with the human OCIF cDNA were isolated using agarose gel electrophoresis and individually subcloned into an EcoRI site of pBluescript II SK + vector (Stratagene) by a conventional method. The resulting plasmids were respectively named pBSE 6, pBSE 4, pBSE 3.6, and pBSE 2.6.

##### (ii) Determination of the nucleotide sequence

The nucleotide sequence of human OCIF genomic DNA which was subcloned into the plasmid was determined using the ABI Dideoxy Terminator Cycle Sequencing Ready Reaction kit (Perkin Elmer) and the 373 Sequencing System (Applied Biosystems). The primer used for the determination of the nucleotide sequence was synthesized based on the nucleotide sequence of human OCIF cDNA (Sequence ID No. 4 in the Sequence Table). The nucleotide

sequences thus determined are given as the Sequences No. 1 and No. 2 in the Sequence Table. The Sequence ID No. 1 includes the first exon of the OCIF gene and the Sequence ID No. 2 includes the second, third, fourth, and fifth exons. A stretch of about 17 kb is present between the first and second exons.

#### Example 4

##### (Production of recombinant OCIF using COS-7 cells)

##### (i) Preparation of OCIF genomic DNA expression cosmid

To express OCIF genomic DNA in animal cells, an expression unit of expression plasmid pCDL-SR $\alpha$ 296 (Molecular and Cellular Biology, vol. 8, P466-472, 1988) was inserted into cosmid vector pWE15 (Stratagene). First of all, the expression plasmid pCDL-SR $\alpha$ 296 was digested with a restriction enzyme Sal I to cut out expression unit with a length of about 1.7 kb which includes an SR promoter, SV40 later splice signal, poly (A) addition signal, and so on. The digestion products were separated by agarose electrophoresis and the 1.7-kb fragment was purified using the QIAEX II gel extraction kit (Qiagen). On the other hand, cosmid vector pWE15 was digested with a restriction enzyme EcoRI and fragments were separated using agarose gel electrophoresis. pWE15 DNA of 8.2 kb long was purified using the QIAEX II gel extraction kit (Qiagen). The ends of these two DNA fragments were blunted using a DNA blunting kit (Takara Shuzo), ligated using a DNA ligation kit (Takara Shuzo), and transferred into *E. coli* DH5 $\alpha$  (Gibco BRL). The resultant transformant was grown and the expression cosmid pWESR $\alpha$  containing an expression unit was purified using a Qiagen column (Qiagen).

The cosmid pWE OCIF containing the OCIF genomic DNA with a length of about 38 kb obtained in (i) above was digested with a restriction enzyme NotI to cut out the OCIF genomic DNA of about 38 kb. After separation by agarose gel electrophoresis, the DNA was purified using the QIAEX II gel extraction kit (Qiagen). On the other hand, the expression cosmid pWESR $\alpha$  was digested with a restriction enzyme EcoRI and the digestion product was extracted with phenol and chloroform, ethanol-precipitated, and dissolved in TE.

pWESR $\alpha$  digested with a restriction enzyme EcoRI and an EcoRI-XmnI-NotI adapter (#1105, #1156 New England Biolaboratory Co.) were ligated using T4 DNA ligase (Takara Shuzo Co., Ltd.). After removal of the free adapter by agarose gel electrophoresis, the product was purified using QIAEX gel extraction kit (Qiagen). The OCIF genomic DNA with a length of about 37 kb which was derived from the digestion with restriction enzyme NotI and the pWESR $\alpha$  to which the adapter was attached were ligated using T4 DNA ligase (Takara Shuzo). The DNA was packaged *in vitro* using the Gigapack packaging extract (Stratagene) and infected with *E. coli* XL1-Blue MR (Stratagene). The resultant transformant was grown and the expression cosmid pWESR $\alpha$ OCIF which contained OCIF genomic DNA was inserted was purified using a Qiagen column (Qiagen). The OCIF expression cosmid pWESR $\alpha$ OCIF was ethanol-precipitated and dissolved in sterile distilled water and used in the following analysis.

##### (ii) Transient expression of OCIF genomic DNA and measurement of OCIF activity

A recombinant OCIF was expressed as described below using the OCIF expression cosmid pWESR $\alpha$ OCIF obtained in (i) above and its activity was measured. COS-7 ( $8 \times 10^5$  cells/well) cells (Riken Cell Bank, RCB0539) were planted in a 6-well plate using DMEM culture medium (Gibco BRL) containing 10% fetal bovine serum (Gibco BRL). On the following day, the culture medium was removed and cells were washed with serum-free DMEM culture medium. The OCIF expression cosmid pWESR $\alpha$ OCIF which had been diluted with OPTI-MEM culture medium (Gibco BRL) was mixed with lipofectamine and the mixture was added to the cells in each well according to the attached protocol. The expression cosmid pWESR $\alpha$  was added to the cells in the same manner as a control. The amount of the cosmid DNA and Lipofectamine was respectively 3  $\mu$ g and 12  $\mu$ l. After 24 hours, the culture medium was removed and 1.5 ml of fresh EX-CELL 301 culture medium (JRH Bioscience) was added to each well. The culture medium was recovered after 48 hours and used as a sample for the measurement of OCIF activity. The measurement of OCIF activity was carried out according to the method described by Kamegawa, M. et al. (Protein, Nucleic Acid, and Enzyme, Vol. 34, p 999 (1989)) and the method of TAKAHASHI, N. et al. (Endocrinology vol. 122, p 1373 (1988)). The osteoclast formation in the presence of activated vitamin D $_3$  from bone marrow cells isolated from mice aged about 17 days was evaluated by the induction of tartaric acid resistant acidic phosphatase activity. The inhibition of the acid phosphatase was measured and used as the activity of the protein which possesses osteoclastogenesis-inhibitory activity (OCIF). Namely, 100  $\mu$ l/well of a OCIF sample which was diluted with  $\alpha$ -MEM culture medium (Gibco BRL) containing  $2 \times 10^{-8}$  M activated vitamin D $_3$  and 10% fetal bovine serum was added to each well of a 96 well micro plate. Then,  $3 \times 10^5$  bone marrow cells isolated from mice (about 17-days old) suspended in 100  $\mu$ l of  $\alpha$ -MEM culture medium containing 10% fetal bovine serum were added to each well of the 96 well micro plate and cultured for a week at 37°C and 100% humidity under 5% CO $_2$  atmosphere. On days 3 and 5, 160  $\mu$ l of the conditioned medium was removed from each well, and 160  $\mu$ l of a sam-

ple which was diluted with  $\alpha$ -MEM culture medium containing  $1 \times 10^{-8}$  M activated vitamin D<sub>3</sub> and 10% fetal bovine serum was added. After 7 days from the start of culturing, the cells were washed with a phosphate buffered saline and fixed with a ethanol/acetone (1:1) solution for one minute at room temperature. The osteoclast formation was detected by staining the cells using an acidic phosphatase activity measurement kit (Acid Phosphatase, Leucocyte, Cat.No. 387-A, Sigma Company). A decrease in the number of cells positive to acidic phosphatase activity in the presence of tartaric acid was taken as the OCIF activity. The results are shown in Table 1, which indicates that the conditioned medium exhibits the similar activity to natural type OCIF obtained from the IMR-90 culture medium and recombinant OCIF produced by CHO cells.

TABLE 1

Activity of OCIF expressed by COS-7 cells in the conditioned medium

Dilution	1/10	1/20	1/40	1/80	1/160	1/320
OCIF genomic DNA introduced	++	++	++	++	+	-
Vector introduced	-	-	-	-	-	-
Untreated	-	-	-	-	-	-

\*++ indicates an activity inhibiting 80% or more of osteoclast formation, \*+ indicates an activity inhibiting 30-80% of osteoclast formation, and - indicates that no inhibition of osteoclast formation is observed.

### (iii) Identification of the product by Western Blotting

A buffer solution (10  $\mu$ l) for SDS-PAGE (0.5 M Tris-HCl, 20% glycerol, 4% SDS, 20  $\mu$ g/ml bromophenol blue, pH 6.8) was added to 10  $\mu$ l of the sample for the measurement of OCIF activity prepared in (ii) above. After boiling for 3 minutes at 100°C, the mixture was subjected to 10% SDS polyacrylamide electrophoresis under non-reducing conditions. The proteins were transferred from the gel to a PVDF membrane (ProBlott, Perkin Elmer) using semi-dry blotting apparatus (Biorad). The membrane was blocked and incubated for 2 hours at 37°C together with a horseradish peroxidase-labeled anti-OCIF antibody obtained by labeling the previously obtained OCIF protein with horseradish peroxidase according to a conventional method. After washing, the protein which has bound the anti-OCIF antibody was detected using the ECL system (Amasham). As shown in Figure 1, two bands, one with a molecular weight of about 120 kilo dalton and the other 60 kilo dalton, were detected in the supernatant obtained from the culture broth of COS-7 cells in which pWESr $\alpha$ OCIF was transfected. On the other hand, these two bands with a molecular weight of about 120 kilo dalton and 60 kilo dalton were not detected in the supernatant obtained from the culture broth of COS-7 cells in which pWESrvector was transfected, confirming that the protein obtained was OCIF.

### INDUSTRIAL APPLICABILITY

The present invention provides a genomic DNA encoding a protein OCIF which possesses an osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA. The protein obtained by expressing the gene of the present invention exhibits an osteoclastogenesis-inhibitory activity and is useful as an agent for the treatment and improvement of diseases involving a decrease in the amount of bone such as osteoporosis, other diseases resulting from bone metabolism abnormality such as rheumatism or degenerative joint disease, and multiple myeloma. The protein is further useful as an antigen to establish antibodies useful for an immunological diagnosis of such diseases.

### NOTE ON MICROORGANISM

#### Depositing Organization:

The Ministry of International Trade and Industry, National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology  
1-3, Higashi-1-Chome, Tsukuba-shi, Ibaraki-ken, Japan  
Date of Deposition: June 21, 1995 (originally deposited on June 21, 1995 and transferred to the international deposition according to the Budapest Treaty on October 25, 1995)  
Accession No. FERM BP-5267

TABLE OF SEQUENCES

Sequence number: 1

Length of sequence: 1316

Sequence Type: nucleic acid

Strandedness: double

Topology: linear

Molecular type: genomic DNA (human OCIF genomic DNA-1)

Sequence:

```

CTGGAGACAT ATAACITGAA CACTTGGCCC TGATGGGGAA GCAGTCTGCG AGGGACTTTT 60
TCAGCCATCT GTAAACAATT TCAGTGGCAA CCGCGAACT GTAATCCATG AATGGACCA 120
CACTTTACAA GTCATCAAGT CTAACITCTA GACCAGCGAA TTAATCGGGG AGACAGCGAA 180
CCCTAGACCA AAGTGCCAAA CTTCGTGCGA TAGCTTGAGG CTAGTGAAA GACCTGAGG 240
AGGCTACTCC AGAAGTTCAG CCGCTAGGAA GCTCCGATAC CAATAGCCCT TTGATGATGG 300
TGGGGTTGGT GAAGGGAACA CTGCTCCGCA AGGTTATCCC TGCCCGAGGC AGTCCAATTT 360
TCACTCTGCA GATTCTCTCT GGCTCTAACT ACCCCAGATA ACAAGCAGTG AATGCAGAAT 420
AGCAGCGGCT TTAGGGCCAA TCAGACATTA GTTAGAAAAA TTCTACTACG ATGGTTTATG 480
TAAACTTGAA GATGAATCAT TCGGAACCTC CCGAAAAGGG CTCACACAAT GCCATGCATA 540
AAGAGGGCCC CTGTAAITTG AGGTTTCAGA ACCCAAGTG AAGGGGTCAG GCAGCCGGGT 600
ACGGCGGAAA CTCACAGCTT TGCCCGACCG AGAGGACAAA GGTCTGGGAC ACACCTCAAC 660
TGCTCCGGA TCTTGGCTGG ATCGGACTCT CAGGCTGGAG GAGACACAAG CACAGCAGCT 720
GCCGAGCGTG TGCCCGACCC TGCCACCGCT GGTCCCGGCT GCCAGGAGCG TGCCCGCTCG 780
CGGGAAGGGG CCGGGAACC TCAGAGCCCG GCGGAGACAG CAGCCGCGCT GTTCTTCAGC 840
CCGCTGGCTT TTTTTCGCC TGCTCTCCCA GGGGACAGAC ACCACCGGCC CACCCTTCAC 900
GCCCACTC CCTGGGGGAT CTTTTCGCC CCAGCCCTCA AACGTTAAT CCTGAGCTT 960
TCTGCACACC CCCCAGCGG TCCCGCCCAA GCTTCTTAAA AAAGAAAGGT GCAAGTTTG 1020
GTCAGGATA GAAAAATCAC TGATCAAAGG CAGGCGATAC TTCCTGTTCG CCGGAGGCTA 1080
TATATAACGT GATGAGCGCA CCGGCTGGGG AGACGCACCG GAGGCTCTCC CCAGCGGCGC 1140

```



CCTCCAAGCC CCTGAGGTTT CCGGGACCA CA ATG AAC AAG TTG CTG TGC TGC 1193

Met Asn Lys Leu Leu Cys Cys

-20

-15

GCG CTC GTG GTAAGTCCCT GGGCCAGCCG ACCGGTCCCC GGGCCCTGGG 1242

Ala Leu Val

GAGGCTGCTG CCACCTGGTG TCCCAACCTC CCAGCGGACC GCGGGGAGA AGGCTCCACT 1302

CGCTCCCTCC CAGG 1316

Sequence number: 2

Length of sequence: 9898

Sequence Type: nucleic acid

Strandedness: double

Topology: linear

Molecular type: genomic DNA (human OCIF genomic DNA-2)

Sequence:

GCTTACTTTG TGCCAAATCT CATTAGGCTT AAGTAATAC AGGACTTTGA GTCAAATGAT 60

ACTGTGGCAC ATAAGAACA ACCTATTTC ATGCTAAGAT GATGCCACTG TGTTCTTTC 120

TCCTTCTAG TTT CTG GAC ATC TCC ATT AAG TGG ACC ACC CAG GAA ACG TTT 171

Phe Leu Asp Ile Ser Ile Lys Trp Thr Thr Gln Gly Thr Phe

-10

-5

1

CCT CCA AAG TAC CTT CAT TAT GAC GAA GAA ACC TCT CAT CAG CTG TTG 219

Pro Pro Lys Tyr Leu His Tyr Asp Glu Glu Thr Ser His Gln Leu Leu

5

10

15

TGT GAC AAA TGT CCT CCT GGT ACC TAC CTA AAA CAA CAC TGT ACA GCA 267

Cys Asp Lys Cys Pro Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala

20 25 30 35

AAG TGG AAG ACC GTG TGC GCC CCT TGC CCT GAC CAC TAC TAC ACA GAC 315

Lys Trp Lys Thr Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp

40 45 50

AGC TGG CAC ACC AGT GAC GAG TGT CTA TAC TGC AGC CCC GTG TGC AAG 363

Ser Trp His Thr Ser Asp Glu Cys Leu Tyr Cys Ser Pro Val Cys Lys

55 60 65

GAG CTG CAG TAC GTC AAG CAG GAG TGC AAT CGC ACC CAC AAC CGC GTG 411

Glu Leu Gln Tyr Val Lys Gln Glu Cys Asn Arg Thr His Asn Arg Val

70 75 80

TGC GAA TGC AAG GAA GGG CGC TAC CTT GAG ATA GAG TTC TGC TTG AAA 459

Cys Glu Cys Lys Glu Gly Arg Tyr Leu Glu Ile Glu Phe Cys Leu Lys

85 90 95

CAT AGG ACC TGC CCT CCT CGA TTT CGA GTG GTG CAA GCT G GTACGTGTCA 509

His Arg Ser Cys Pro Pro Gly Phe Gly Val Val Gln Ala

100 105 110

ATGTGCAGCA AAATTAATTA GGATCATGCA AAGTCAGATA GTTGTGACAG TTTAGGAGAA 569

CACTTTTGT CTGATGACAT TATAGGATAG CAAATTGCAA AGGTAATGAA ACCTGCCAGG 629  
 5 TAGTACTAT GTGTCTGGAG TGCTTCCAAA GGACCATTGC TCAGAGGAAT ACTTTGCCAC 689  
 TACAGGGCAA TTTAATGACA AATCTCAAAAT GCAGCAAAAT ATTCTCTCAT GAGATCCAT 749  
 ATGTTTTTTT TTTTTTTTTT TAAAGAAACA AACTCAAGTT GCACTATTGA TAGTTGATCT 809  
 10 ATACCTCTAT ATTTCACTTC AGCATGGACA CCTTCAAACT GCAGCAGTTT TTGACAAACA 869  
 TCAGAAATGT TAATTATAC CAAGAGAGTA ATTATGCTCA TATTAATCAG ACTCTGCAGT 929  
 15 GCTAACATA AGCAGTTATA ATTAATTATG TAAAAAATGA GAATGGTCAG CGGAATTGCA 989  
 TTTCATTATT AAAACAAGG CTAGTTCTTC CTTTAGCATG GGAGCTGAGT GTTTGGGAGG 1049  
 GTAAGGACTA TAGCAGAATC TCTTCAATGA GCTTATTCTT TATCTTAGAC AAAACAGATT 1109  
 20 GTCAGGCCAA GAGCAAGCAC TTGECTATAA ACCAAGTGCT TTCTCTTTTG CATTTTGAAC 1169  
 AGCATTTGTC AGGGCTCATG TGTATTGAAT CTTTAAACC AGTAACCCAC GTTTTTTTTC 1229  
 TGCCACATTT CGGAAGCTTC AGTGCAGCCT ATAACTTTTC ATAGCTTGAG AAAATTAAGA 1289  
 25 GTATCCACTT ACTTAGATCG AAGAAGTAAT CAGTATAGAT TCTGATGACT CAGTTTGAAC 1349  
 CAGTGTITCT CAACTGAAGC CCTGCTGATA TTTTAAGAAA TATCTGGATT CCTAGGCTCG 1409  
 30 ACTCCTTTTT GTGGGAGCT GTCTGCGCA TTGTAGAATT TTGCGAGCAC CCCTGGACTC 1469  
 TAGCCACTAG ATACCAATAG CAGTCCTTCC CCCATGTGAC AGGCCAAAAAT GTCTTCAGAC 1529  
 ACTGTCAAAT GTGGCCAGCT GGCAAAATCA CTCCTGGTTC AGAACAGGCT CATCAATGCT 1589  
 35 AAGTATCTGT AACTATTTTA ACTCTCAAAA CTGTGTGATAT ACAAAGTCTA AATTATTAGA 1649  
 CGACCAATAC TTTAGGTTTA AAGGCATACA AATGAAACAT TCAAAAATCA AAATCTATTG 1709  
 40 TGTTTCTCAA ATAGTGAATC TTATAAAAT AATCAGAGAA CATGCAAAAT GCATCAGACT 1769  
 CCTTAAAT TCCTCTTCGT ATGAGTATTG GAGGAGGAA TTGGTGATAG TTCTACTT 1829  
 45 CTATTGGATG GTACTTTGAG ACTCAAAAGC TAAGCTAAGT TGTGTGTGTG TCAGGGTGGC 1889  
 GGGTGTGGAA TCCCATCAGA TAAAGCAAA TCCATGTAAT TCATTGAGTA AGTTGTATAT 1949  
 GTAGAAAAAT GAAAAGTGGG CTATGCAGCT TCGAACTAG AGAATTTTGA AAAAATATGC 2009  
 50 AAATCACAAG GATCTTTCTT AAATAAGTAA GAAATCTGT TTGTAGAATC AAGCAAGCAG 2069  
 GCAGCCAGAA GACTCAGAAC AAAAGTACAC ATTTTACTCT GTGTACACTG GCAGCAGACT 2129

GGGATTTATT TACCTCTCCC TCCCTAAAAA CCCACACAGC GGTTCCTCTT GCGAAATAAG 2189  
 5 AGGTTTCCAG CCCAAAGAGA AGGAAAGACT ATGTGGTGTT ACTCTAAAAA GTATTTAATA 2249  
 ACCGTTTTGT TGTGCTGTT GCTGTTTTGA AATCAGATTG TCTCCTCTCC ATATTTTATT 2309  
 TACTTCATTC TGTTAATTCC TGTGGAATTA CTTAGAGCAA GCATGGTGAA TTCTCAACTG 2369  
 10 TAAAGCCAAA TTTCTCCATC ATTATAATTT CACATTTTGC CTGGCAGGTT ATAAATTTTA 2429  
 TATTTCCACT GATAGTAATA AGGTAAAATC ATTACTTAGA TGGATAGATC TTTTTCATAA 2489  
 15 AAAGTACCAT CAGTTATAGA GGAAGTCAT GTTCATGTTT AGGAAGGTCA TTAGATAAAG 2549  
 CTTCTGAATA TATTATGAAA CATTAGTTCT GTCAATTCTTA GATTCITTTT GTTAAATAAC 2609  
 TTTAAAAGCT AACTTACCTA AAAGAAATAT CTGACACATA TGAACCTCTC ATTAGGATGC 2669  
 20 AGGAGAAGAC CCAAGCCACA GATATGTATC TGAAGAATGA ACAAGATTCT TAGGCCCGGC 2729  
 ACGGTGCGTC ACATCTGTAA TCTCAAGACT TTGAGAGGTC AAGGCGGGCA GATCACCTGA 2789  
 25 GGTGAGGAGT TCAAGACCAG CCTGGCCAAC ATGATGAAAC CTTGCTCTTA CTAATAATAC 2849  
 AAAAATTAGC AGGGCATGCT GGTGCATGCC TGCAACCCTA GCTACTCAGG AGGCTGAGAC 2909  
 ACGAGAATCT CTGAAACCTC CGAGGCGGAG GTTGTGGTGA GCTGAGATCC CTCTACTGCA 2969  
 30 CTCCAGCGCT GGTGACAGAG ATGAGACTCC GTCCCTGCCG CCGCCCCCGC CTTCCCCCCC 3029  
 AAAAAGATTC TTCTTCATGC AGAACATACG GCAGTCAACA AAGGGAGACC TGGGTCCAGG 3089  
 TGTCCAAGTC ACTTATTTGG ACTAAATTAG CAATGAAAGA ATGCCATGGA ATCCCTGCCC 3149  
 35 AAATACCTCT GCTTATGATA TTGTAGAATT TGATATAGAG TTGTATCCCA TTTAAGGAGT 3209  
 AGGATGTAGT AGGAAAGTAC TAAAAACAAA CACACAACA GAAAACCCCTC TTTGCTTTGT 3269  
 40 AAGGTGGTTC CTAAGATAAT GTCAGTGCAA TGCTGGAAT AATATTTAAT ATGTGAAGGT 3329  
 TTTAGGCTGT GTTTTCCCTC CTGTTCTTTT TTTTCTGCCA GCCCTTTGTC ATTTTTCGAG 3389  
 45 GTCAATGAAT CATGTAGAAA GAGACAGGAG ATGAAACTAG AACCACTCCA TTTTGCCCTC 3449  
 TTTTATATTT TCTGGTTTTG GTAAAAGATA CAATGAGGTA GGAGGTTGAG ATTTATAAAT 3509  
 GAAGTTTAAT AAGTTTCTGT AGCTTTGATT TTTCCTTTTC ATATTGTGTA TCTTGCAATA 3569  
 50 GCCAGAATTG GCCTGTAAAA TCTACATATG GATATTGAAG TCTAAATCTG TTCAACTAGC 3629  
 TTACACTAGA TGGAGATATT TTCATATTCA GATACACTGG AATGTATGAT CTAGCCCATG 3689

5 GTAAATAGT CAAGTGTTG AAGGTATTIA TTTTAAATAG CGTCTTAGT TGTGGACTGG 3749  
 TTCAAGTTTT TCTGCCAATG ATTCTTCAA AITTATCAAA TATTTTCCA TCATGAAGTA 3809  
 AAATGCCCTT GCAGTCACCC TTCTGAAGT TTGAACGACT CTGCTGTTT AAACAGTTTA 3869  
 10 ACCAAATGGT ATATCATCTT CCGTTTACTA TGTAGCTTAA CTGCAGGCTT ACGCTTTTGA 3929  
 GTCAGCGGCC AACTTTATTG CCACCTTCAA AAGTTTATTA TAATGTTGA AATTTTACT 3989  
 TCTCAAGGT AGCATACTTA GGAGTTGCTT CACAATTAGG ATTCAGCAA GAAAGAAGTT 4049  
 15 CAGTAGGAAC TGATTGGAAT TTAATGATGC AGCATTCAAT GGTACTAAT TTCAAAGAAT 4109  
 GATATTACAG CAGACACACA GCAGTTATCT TGATTTTCTA GGAATAATTG TATGAAGAAT 4169  
 ATGGTGACA ACACGGGCTT ACTGCCACTC AGCGGAGCTT GGACTAATGA ACACCTTACC 4229  
 20 CTCTTTCTT TTCTCTCAC ATTTCATGAG CGTTTTGTAG GTAACGAGAA AATTGACTTG 4289  
 CATTGCGATT ACAAGGAGGA GAAACTGGCA AAGGGGATGA TGTGCGAAGT TTGTGCTGT 4349  
 25 CTAATCAAGT GAAAAATGAA AATGCTAGAG TTTTGTGCAA CATAATAGTA GCAGTAAAAA 4409  
 CCAAGTGAAA AGTCTTTCCA AAAGTGCTT AAGAGGGCAT CTGCTGGGAA ACGATTTCAG 4469  
 GAGAAGGTAC TAAATTGCTT GGTATTTTCC GTAG GA ACC CCA GAG CGA AAT ACA 4523

Gly Thr Pro Glu Arg Asn Thr

115

35 GTT TGC AAA AGA TGT CCA GAT GCG TTC TTC TCA AAT CAG ACG TCA TCT 4571  
 Val Cys Lys Arg Cys Pro Asp Gly Phe Phe Ser Asn Glu Thr Ser Ser  
 40 120 125 130 135

45 AAA CCA CCC TGT AGA AAA CAC ACA AAT TGC AGT GTC TTT GGT CTC CTG 4619  
 Lys Ala Pro Cys Arg Lys His Thr Asn Cys Ser Val Phe Gly Leu Leu  
 50 140 145 150

CTA ACT CAG AAA GGA AAT GCA ACA CAC GAC AAC ATA TGT TCC GGA AAC 4687

Leu Thr Glu Lys Gly Asn Ala Thr His Asp Asn Ile Cys Ser Gly Asn

155

160

165

AGT GAA TCA ACT CAA AAA TGT GGA ATA G GTAATTACAT TCCAAAATAC 4715

Ser Glu Ser Thr Glu Lys Cys Gly Ile

170

175

GCTTTGTAC GATTTGTAG TATCATCTCT CTCTCTGAGT TGAACACAAG GCCTCCAGCC 4775  
 ACATTCTTGG TCAAACTTAC ATTTTCCCTT TCTTGAATCT TAACCAGCTA AGGCTACTCT 4835  
 CGATGCATTA CTGCTAAAGC TACCACTCAG AATCTCTCAA AAACCTCATCT TCTCAGAGAT 4895  
 AACACCTCAA AGCTTGATTT TCTCTCCTTT CACACTGAAA TCAAATCTTG CCGATAGGCA 4955  
 AAGGGCAGTG TCAAGTTTGC CACTGAGATG AAATTAGGAG AGTCCAAACT GTAGAATTCA 5015  
 CGTTGTGTGT TATTACTTTC ACGAATGTCT GTATTATTAA CTAAAGTATA TATTGGCAAC 5075  
 TAAGAAGCAA AGTGATATAA ACATGATGAC AAATTAGGCC AGGCATGGTG GCTTACTCCT 5135  
 ATAATCCCAA CATTTTGGGG GCCCAAGGTA GGCAGATCAC TTGAGGTGAG GATTTCAAGA 5195  
 CCAGCCTGAC CAACATGGTG AAACCTTGTC TCTACTAAAA ATACAAAAAT TAGCTGGCCA 5255  
 TGGTAGCAGG CACTTCTAGT ACCAGCTACT CAGGGCTGAG GCAGGAGAAT CGCTTGAACC 5315  
 CAGGAGATGG AGGTTGCAGT GAGCTGAGAT TGTACCACTG CACTCCAGTC TGGGCAACAG 5375  
 AGCAAGATTT CATCACACAC ACACACACAC ACACACACAC ACACATTAGA AATGTGTACT 5435  
 TGGCTTTGTT ACCTATGCTA TTAGTGCATC TATTGCATGG AACTTCCAAG CTACTTCGCT 5495  
 TGTGTTAAGC TCTTCATTGG GTACAGGTCA CTAGTATTAA GTTCAGGTTA TTCGGATGCA 5555  
 TTCCACGGTA GTGATGACAA TTCATCAGGC TAGTGTGTGT GTTCACCTTG TCACTCCAC 5615  
 CACTAGACTA ATCTCAGACC TTCACTCAA GACACATTAC ACTAAGATG ATTTGCTTTT 5675  
 TTGTGTTTAA TCAAGCAATG GTATAACCA GCTTGACTCT CCCCAAACAG TTTTGTGATC 5735  
 TACAAAGAAG TTTATGAAGC AGAGAAATGT GAATTGATAT ATATATGAGA TTCTAACCCA 5795  
 GTTCCAGCAT TGTTTCATTG TGTAAITGAA ATCATAGACA AGCCATTITTA GCCTTTGCTT 5855

TCTTATCTAA AAAAAAAAAA AAAAAATGA AGCAAGCGGT ATTAAAGGA GTGATCAAT 5915  
 5 TTAAACATTC TCTTAAATTA ATTCAITTTT AATTTTACTT TTTTTCATT ATTGTGCACT 5975  
 TACTATGTGG TACTGTGCTA TAGAGGCTTT AACATTATA AAAACACTGT GAAAGTTGCT 6035  
 TCAGATGAAT ATAGGTAGTA GAACGGCAGA ACTAGTATTC AAAGCCAGGT CTGATGAATC 6095  
 10 CAAAAACAAA CACCCATTAC TCCCATTTTC TGGGACATAC TTACTCTACC CAGATCTCT 6155  
 GGGCTTTGTA ATGCCATGCT AAATAACATA GTTTTATGTT TGGTTATTTT CCTATGTAAT 6215  
 15 GTCTACTTAT ATATCTGTAT CTATCTCTTG CTITGTTTCC AAAGGTAAC TATGTGCTA 6275  
 AATGTGGGCA AAAAAAACA CACTATTCCA AATTACTGTT CAAATTCCTT TAAGTCAGTG 6335  
 ATAATTAATT GTTTTGACAT TAATCATGAA GTTCCCTGTG GCTACTAGGT AAACCTTAA 6395  
 20 TAGAATGTTA ATGTTTGAT TCAATTAAAG AATTTTGGC TGTACTTAT TTACAACAAT 6455  
 AATTCACCTT AATTAGACAT TTAATAAAT TTCTCTTCAA AACATGCCC AAAAAAGAAC 6515  
 ATTAGAAGAC ACCTAAGCTC AGTTGGTCTC TGCCACTAAG ACCAGCCAAC AGAAGCTTGA 6575  
 25 TTTTATTCAA ACTTTGCATT TTAGCATATT TTATCTTGA AATTTCAATT GTGTGGTTT 6635  
 TTTGTTTTG TTTGTATTGA ATAGACTCTC AGAAATCCAA TTGTGACTA AATCTCTCG 6695  
 30 GTTTTCTAAC CTTCTTTAG AT GTT ACC CTG TGT GAG GAG GCA TTC TTC AGG 6747

Asp Val Thr Leu Cys Glu Glu Ala Phe Phe Arg

180

185

35

TTT GCT GTT CCT ACA AAG TTT ACG CCT AAC TGG CTT AGT GTC TTG GTA 6795  
 40 Phe Ala Val Pro Thr Lys Phe Thr Pro Asn Trp Leu Ser Val Leu Val  
 190 195 200

45

GAC AAT TTG CCT GGC ACC AAA GTA AAC GCA GAG AGT GTA GAG AGG ATA 6843  
 Asp Asn Leu Pro Gly Thr Lys Val Asn Ala Glu Ser Val Glu Arg Ile  
 50 205 210 215

55

AAA CGG CAA CAC AGC TCA CAA GAA CAG ACT TTC CAG CTC CTC AAG TTA 6891

Lys Arg Gln His Ser Ser Gln Glu Gln Thr Phe Gln Leu Leu Lys Leu

220 225 230 235

TGG AAA CAT CAA AAC AAA GAC CAA GAT ATA GTC AAG AAG ATC ATC CAA G 6940

Trp Lys His Gln Asn Lys Asp Gln Asp Ile Val Lys Lys Ile Ile Gln

240 245 250

GTATGATAAT CTAATAAAA AAGATCAATC AGAAATCAAA GACACCTATT TATCATAAAC 7000

CAGGAACAAG ACTGCATGTA TGTTTAGTTG TGTGGATCTT GTTCCCTGTG TGAATCATT 7060

GTTGGACTGA AAAAGTTTCC ACCTGATAAT GTAGATGTGA TTCCACAAC AGTTATACAA 7120

GGTTTTGTTC TCACCCCTGC TCCCAGTTT CCTTGTAAG TATGTTGAAC ACTCTAAGAG 7180

AAGAGAAATG CAITTTAAGG CAGGGCTGTA TCTCAGGGAG TCGCTTCCAG ATCCCTTAAC 7240

GCTTCTGTAA GCAGCCCTC TAGACCACCA AGGAGAAGCT CTATAACCAC TTGTATCTT 7300

ACATTGCACC TCTACCAAGA AGCTCTGTTG TATTTACTTG GTAATTTCTT CCAGGTAGGC 7360

TTTTCGTAGC TTACAAATAT GTTCTTATTA ATCTCATGA TATGGCCTGC ATTAATAATTA 7420

TTTTAATGCC ATATGTTATG AGAATTAATC AGATAAAATC TGAAGCTGT TTGAGCCTCT 7480

TGTAGGAAAA ACCTAGTTAC AGCAAAATGT TCTCACATCT TATAAGTTTA TATAAAGATT 7540

CTCCTTTAGA AATGGTGTGA GAGAGAAACA GAGAGAGATA GGGAGAGAAG TGTGAAGAA 7600

TCTGAAGAAA AGGAGTTTCA TCCAGTGTGG ACTGTAAGCT TTACGACACA TGATGGAAG 7660

AGTTCTGACT TCAGTAAGCA TTGGGAGGAC ATGCTAGAAG AAAAAGGAAG AAGAGTTTCC 7720

ATAATGCAGA CAGGTCAGT GAGAAATTCA TTCAGGTCCT CACCAGTAGT TAAATGACTG 7780

TATAGTCTTG CACTACCTTA AAAAATCTCA AGTATCTGAA ACCGGGCCAA CAGATTTTAG 7840

GAGACCAACG TCTTTGAGAG CTGATTGCTT TTGCTTATGC AAAGAGTAAA CTTTATGTT 7900

TTGAGCAAC CAAAGTATT CTTTGAACGT ATAATTAGCC CTGAAGCCGA AAGAAAAGAG 7960

AAAATCAGAG ACCGTTAGAA TTGGAAGCAA CCAAAATCCC TATTTTATAA ATGAGGACAT 8020



5 TTTAACCAG AAAGATGAAC CGATTGGCT TAGGGCTCAC AGATACTAAG TGACTCATGT 8080  
 CATTAATAGA AATGTTAGTT CCTCCCTCTT AGGTTTGTA CCTAGCTTAT TACTGAAATA 8140  
 TTCTTAGGC TGTGTGCTC CTTTAGTCC TCGACCTCAT GTCTTTGAGT TTTCAGATAT 8200  
 CCTCCTCATG GAGGTAGTCC TGTGGTGCTA TGTGTATTCT TTAAGGCTA GTTACGGCAA 8260  
 10 TTAAGTATC AACTAGCGCC TACTAATGAA ACTTTGTATT ACAAGTAGC TAACTTGAAT 8320  
 ACITTCCTTT TTTTCTGAAA TGTATGGTG GTAATTCTC AAACITTTTC TTAGAAAAC 8380  
 15 GAGAGTCATG TGTCTTATT TCTACTGTA ATTTCAAAA TTAGGAGCTT CTCCAAAGT 8440  
 TTTGTTGGAT CCCAAAAATA TATAGCATAT TATCTTATTA TAACAAAAA TATTTATCTC 8500  
 AGTTCCTAGA AATAAATGGT GTCACCTAAC TCCCTCTCAA AAGAAAAGG TATCATTGAA 8560  
 20 ATATAATTAT GAAATTCGC AAGAACCCTT TGCCACGC TTGTTTATG ATGGCATTGG 8620  
 ATGAATATAA ATGATGTGAA CACTTATCTG GCGTTTCTCT TTATGCAG AT ATT GAC 8676

Asp Ile Asp

25  
 30 CTC TGT GAA AAC AGC GTG CAG CGG CAC ATT GGA CAT GCT AAC CTC ACC 8724  
 Leu Cys Glu Asn Ser Val Gln Arg His Ile Gly His Ala Asn Leu Thr  
 255 260 265 270

35  
 40 TTC GAG CAG CTT CGT AGC TTG ATG GAA AGC TTA CCG GGA AAG AAA GTG 8772  
 Phe Glu Gln Leu Arg Ser Leu Met Glu Ser Leu Pro Gly Lys Lys Val  
 275 280 285

45  
 50 GGA GCA GAA GAC ATT GAA AAA ACA ATA AAG GCA TGC AAA CCC AGT GAC 8820  
 Gly Ala Glu Asp Ile Glu Lys Thr Ile Lys Ala Cys Lys Pro Ser Asp  
 290 295 300

55  
 CAG ATC CTG AAG CTG CTC AGT TTG TGG CGA ATA AAA AAT GGC GAC CAA 8868

Gln Ile Leu Lys Leu Leu Ser Leu Trp Arg Ile Lys Asn Gly Asp Gln  
 305 310 315

GAC ACC TTC AAG GGC CTA ATG CAC GCA CTA AAG CAC TCA AAG ACC TAC 8916  
 Asp Thr Leu Lys Gly Leu Met His Ala Leu Lys His Ser Lys Thr Tyr  
 320 325 330

CAC TTT CCC AAA ACT GTC ACT CAG AGT CTA AAG AAG ACC ATC AGG TTC 8964  
 His Phe Pro Lys Thr Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe  
 335 340 345 350

CTT CAC AGC TTC ACA ATG TAC AAA TTG TAT CAG AAG TTA TTT TTA GAA 9012  
 Leu His Ser Phe Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu  
 355 360 365

ATG ATA GGT AAC CAG GTC CAA TCA GTA AAA ATA AGC TGC TTA 9054  
 Met Ile Gly Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu  
 370 375 380

TAAGTGGAAA TGGCCATTGA GCTGTTTCCT CACAATTGGC GAGATCCCAT GGATGACTAA 9114  
 ACTGTTTCTC AGGCACTTGA GGCTTTCAGT GATATCTTTC TCATTACCAG TCACTAAITT 9174  
 TGCCACAGGG TACTAAAAGA AACTATGATG TGGAGAAAGG ACTAACATCT CTCCAATAA 9234  
 ACCCCAAATG GTTAATCCAA CTGTCAGATC TGGATCGTTA TCTACTGACT ATATTTTCCC 9294  
 TTATTACTGC TTGCACTAAT TCAACTGGAA ATTAACAAAA AAAAAGTAGA CTCCACTGGG 9354  
 CCTTACTAAA TATGGGAATG TCTAACTTAA ATAGCTTTGG GATTCCAGCT ATGCTAGAGG 9414  
 CTTTATTAG AAAGCCATAT TTTTTCCTGT AAAAGTTACT AATATATCTG TAACACTATT 9474

ACAGTATTGC TATTATATT CATTGAGATA TAAGATTGG ACATATTATC ATCCTATAAA 9534  
 5 GAAACGGTAT GACTTAATT TAGAAAGAAA ATTATATTCT GTTATTATG ACAAATGAAA 9594  
 GAGAAAATAT ATATTTTIAA TGGAAACTTT GTAGCATTTT TCTAATAGGT ACTGCCATAT 9654  
 TTTTCTGTGT GGAGTATTTT TATAATTTTA TCTGTATAAG CTGTAATATC ATTTTATAGA 9714  
 10 AAATGCATTA TTTAGTCAAT TGTTTAATGT TGGAAAACAT ATGAAATATA AATTATCTGA 9774  
 ATATTAGATG CTCTGAGAAA TTGAAATGAC CTTATTTAAA AGATTTTATG GTTTTATAAC 9834  
 TATATAAATG ACATTATTAA AGTTTTCAAA TTATTTTITA TTGCTTTCTC TGTTCCTTTT 9894  
 15 ATTT 9898

Sequence number: 3

Length of sequence: 401

Sequence Type: amino acid

Strandedness: single stranded

Topology: linear

Molecular type: protein

Sequence:

Met Asn Asn Leu Leu Cys Cys Ala Leu Val Phe Leu Asp Ile Ser

-20

-15

-10

Ile Lys Trp Thr Thr Gln Glu Thr Phe Pro Pro Lys Tyr Leu His

-5

1

5

Tyr Asp Glu Glu Thr Ser His Gln Leu Leu Cys Asp Lys Cys Pro

10

15

20

Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala Lys Trp Lys Thr

25

30

35

Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp Ser Trp His

40

45

50

Ile Gln Asp Ile Asp Leu Cys Glu Asn Ser Val Gln Arg His Ile

250 255 260

Gly His Ala Asn Leu Thr Phe Glu Gln Leu Arg Ser Leu Met Glu

265 270 275

Ser Leu Pro Gly Lys Lys Val Gly Ala Glu Asp Ile Glu Lys Thr

280 285 290

Ile Lys Ala Cys Lys Pro Ser Asp Gln Ile Leu Lys Leu Leu Ser

295 300 305

Leu Trp Arg Ile Lys Asn Gly Asp Gln Asp Thr Leu Lys Gly Leu

310 315 320

Met His Ala Leu Lys His Ser Lys Thr Tyr His Phe Pro Lys Thr

325 330 335

Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe Leu His Ser Phe

340 345 350

Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu Met Ile Gly

355 360 365

Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu

370 375 380

Sequence number: 4

Length of sequence: 1206

Sequence Type: nucleic acid

Strandedness: single stranded

Topology: linear

Molecular type: cDNA

## Sequence:

5 ATGAACAACT TGCTGTGCTG CGCCTCGTG TTTCTGGACA TCCTCATTAA GTGACCACC 60  
 CAGGAAACGT TTCTCCAAA GTACCTTCAT TATGACGAAG AAACCTCTCA TCAGCTGTTG 120  
 TGTGACAAAT GTCTCCTGG TACCTACCTA AAACAACACT GTACAGCAAA GTGGAAGACC 180  
 10 GTGTGCGCCC CTGCCCCTGA CCACTACTAC ACAGACAGCT GGCACACCAG TGACGAGTGT 240  
 CTATACTGCA GCCCCTGTG CAAGGAGCTG CAGTACGTCA AGCAGGAGTG CAATCCACC 300  
 15 CACAACCCGG TGTGCGAATG CAAGGAAGGG CGCTACCTTG AGATAGAGTT CTGCTTGAAA 360  
 CATAGGAGCT GCCCTCCTGG AITTTGGAGTG GTGCAAGCTG GAACCCCAAG GCGAAATACA 420  
 GTTTGCAAAA GATGTCCAGA TGGGTTCTTC TCAAATGAGA CGTCATCTAA AGCACCCCTGT 480  
 20 AGAAAACACA CAAATTGCAG TGTCTTTGGT CTCTGTCTAA CTCAGAAAGG AAATGCAACA 540  
 CAGGACAACA TATGTTCCGG AAACAGTGAA TCAACTCAAA AATGTGGAAT AGATGTTACC 600  
 25 CTGTGTGAGG AGGCATTCTT CAGGTTTGCT GTTCTACAAA AGTTTACCCC TAACTGGCTT 660  
 AGTGTCTTGG TAGACAATTT GCCTGGCACC AAAGTAAACG CAGAGAGTGT AGAGAGGATA 720  
 AAACGGCAAC ACAGCTCACA AGAACAGACT TTCCAGCTGC TGAAGTTATG GAAACATCAA 780  
 30 AACAAGACC AAGATATAGT CAAGAAGATC ATCCAAGATA TTGACCTCTG TGAACACAGC 840  
 GTCCAGCGGC ACAITGGACA TGCTAACCTC ACCTTCGAGC AGCTTCCTAG CTTGATCGAA 900  
 35 AGCTTACCGC GAAAGAAAGT GCGAGCAGAA GACATTGAAA AAACAATAAA GGCATGCAAA 960  
 CCCAGTGACC AGATCTTGAA GCTGCTCAGT TTGTGGCGAA TAAAAAATGG CGACCAAGAC 1020  
 ACCTTGAAGG GCCTAATGCA CCACTAAAG CACTCAAGA CGTACCACTT TCCCAAACT 1080  
 40 GTCACCTAGA GTCTAAAGAA GACCATCAGG TTCCTTCACA GCTTCACAAT GTACAAATTG 1140  
 TATCAGAAGT TATTTTTAGA AATGATAGGT AACCAAGTCC AATCAGTAAA AATAAGCTGC 1200  
 45 TTATAA 1206

50

55

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: SNOW BRAND MILK PRODUCTS CO., LTD.  
 (B) STREET: 1-1, MAEBOCHO 6-CHOME  
 (C) CITY: NIGASHI-KU, SAPPORO-SHI  
 (D) STATE: HOKKAIDO  
 (E) COUNTRY: JP  
 (F) POSTAL CODE (ZIP): NONE

## (ii) TITLE OF INVENTION: NOVEL DNA AND PROCESS FOR PREPARING PROTEIN USING THE DNA

## (iii) NUMBER OF SEQUENCES: 4

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
 (B) COMPUTER: IBM PC compatible  
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS  
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

## (v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: EP 97935810.8

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: JP 235928/96  
 (B) FILING DATE: 19-AUG-1996

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1316 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: genomic DNA (human OCIF genomic DNA-1)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTGGAGACAT ATAACCTGAA CACTTGGCCC TGATGGGGAA GCAGCTCTGC AGGGACTTTT 60  
 TCAGCCATCT GTAACAAT TCAGTGGCAA CGCGCGAACT GTAATCCATG AATGGGACCA 120  
 CACTTTACAA GTCAACAAGT CTAACCTCTA GACACGGGAA TTAATGGGGG ACACAGCGAA 180  
 CCTTAGACAA AAGTGCCAAA CTCTCTGCGA TAGCTTTAGG CTAGTGAA GACTCCAGG 240  
 AGGCTACTCC AAGAGTTCAG CGCTAGGAA GCTCCGATAC CAATAGCCCT TTGATGATGG 300  
 TGGGTTGGT GAAGGGAACA GTGCTCCGCA AGGTTATCC TCCCCAGGC AGTCCATT 360  
 TCAGTCTGCA GATTCCTCT GGCCTAACT ACCCCAGATA ACAAGGAGTG AATCCAGAA 420  
 AGCAGGGCT TTAGGGCCAA TCAGACATTA GTTAGAAAA TTCTACTAC ATGGTTTATG 480  
 TAAACTTGAA GATGAATAT TCGCAACTC CCGAAAGGG CTCAGACAA GCATCATATA 540  
 AAGAGGGGCC CTGTAATTG AGGTTTCAAG ACCCGATGT AAGGGGTGAG GCAGCGGGGT 600  
 ACGCGGGAAA CTCACAGCTT TGGCCACGG AGAGGACAAA GGTCTGGAC ACATCCAC 660  
 TGGCTCCGGA TTTGGCTGG ATCGGACTCT CAGGGTGGAG GAGACACAAG CACAGCAGCT 720  
 GCCAGCGTG TGCACAGCCC TCCACCGCT GGTCCGGCT GCCAGGAGGC TGGCGGCTGG 780  
 CGGAGGGGG CCGGAAACC TCAGAGCCCC GGGGAGACAG CAGCGGCTT GTTCCTCAGC 840  
 CGGGGCTT TTTTCCCGT TGCTCTCCA GGGGACAGAC ACCACCGGCC CACCGCTCAC 900  
 CGCCACCTC CCGGGGGAT CTCTTCCA CAGCGCTTAA AGGCTTAAT CTGGAGCTT 960  
 TCTGACACC CCGCCACCG TCCCGCCAA GAGCGGATAC TTCTCTTTC CCGGAGCTA 1020  
 GTCCAGGATA GAAAAATGAC TGATCAAGG CAGCGGATAC TTCTCTTTC CCGGAGCTA 1080  
 TATATAACT GATGAGCGCA CGGGCTGGG AGAGCCAGCG GAGCGCTGC CCGCGCGCG 1140  
 CCTCCAGCC CTTGAGTTT CCGGGGACCA CA ATG AAC AAG TTG CTG TGC TGC 1193  
 Met Asn Lys Leu Cys Cys  
 -20 -15

GCG CTC GTG GTAATCCCT GGGCAGCGG ACGGGTGGCC GCGCGCTGG 1242

Ala Leu Val

GAGGCTGCTG CCACCTGGTC TCCCAACCTC CCAGCGGACC GCGCGGGAGA AGGCTCCACT 1302  
 CGTCCCTCC CAGG 1316

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 989 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic DNA (human OCIF genomic DNA-2)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GCTTACTTTG TCCAAATCT CATTAGGCTT AAGGTAATAC AGGACTTTGA GTCAAATGAT 60  
 ACTGTTGCAC ATAAGAACAA ACCTATTTC ATGCTAAGAT GATGCCACTG TGTCTCTTC 120  
 TCTCTCTAG TTT CTG GAC ATC TCC ATT AAG TGG ACC ACC CAG GAA ACG TTT 171  
 Phe Leu Asp Ile Ser Ile Lys Trp Thr Thr Gln Glu Thr Phe  
 -10 -5 1  
 OCT CCA AAG TAC CTT CAT TAT GAC GAA GAA ACC TCT CAT CAG CTG TTG 219  
 Pro Pro Lys Tyr Leu His Tyr Asp Glu Glu Thr Ser His Gln Leu Leu  
 5 10 15  
 TGT GAC AAA TOT CCT CCT GGT ACC TAC CTA AAA CAA CAC TGT ACA GCA 267  
 Cys Asp Lys Cys Pro Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala  
 20 25 30  
 AAG TGG AAG ACC GTG TGC GCC CCT TGC CCT GAC CAC TAC TAC ACA GAC 315  
 Lys Trp Lys Thr Val Cys Ala Pro Cys Pro Asp His Tyr Thr Asp  
 40 45 50  
 AGC TGG CAC ACC AGT GAC GAG TGT CTA TAC TGC AGC CCC GTG TGC AAG 363  
 Ser Trp His Thr Ser Asp Glu Cys Leu Tyr Cys Ser Pro Val Cys Lys  
 55 60 65  
 GAG CTG CAG TAC GTC AAG CAG GAG TGC AAT CGC ACC CAC AAC CGC GTG 411  
 Glu Leu Gln Tyr Val Lys Gln Glu Cys Asn Arg Thr His Asn Arg Val  
 70 75 80  
 TGC GAA TGC AAG GAA GGG CGC TAC CTT GAG ATA GAG TTC TGC TTG AAA 459  
 Cys Glu Cys Lys Glu Gly Arg Tyr Leu Glu Ile Glu Phe Cys Leu Lys  
 85 90 95  
 CAT AGG AGC TGC CCT CCT GGA TTT GGA GTG GTG CAA GCT G GTACGTGTCA 509  
 His Arg Ser Cys Pro Pro Gly Phe Gly Val Val Gln Ala  
 100 105 110  
 ATGTGCAGCA AAATTANTTA GGATCATGCA AAGTCAGATA GTTGTGACAG TTTAGGAGAA 569  
 CACTTTTGT CTGATGACAT TATAGGATAG CAAATTGCAA AGGTAATGAA ACCTGCCAGG 629  
 TAGTACTAT GTGTCTGGAG TGCTTCCAAA GGACCATTGC TCAGAGGAAT ACTTTGCCAC 689  
 TCAGGGGCAA TTAAATGACA AATCTCAAA GTACGCAAAAT ATTCTCTCAT GAGATGCAATG 749  
 ATGGTTTTT TTTTTTTTT TAAAGAAACA AACTCAAGTT GCACATATGA TAGTGATCT 809  
 ATACCTCTAT ATTTCACCTC AGCATGGACA CTTCTAACCT GCAGCACTTT TTGACAAACA 869  
 TCAGAAATGT TAATTATAC CAAGAGASTA ATTATGCTCA TATTAATGAG ACTCTGGAGT 929  
 CTATACATA AGCAATTATA ATTAATTATG TAAAGAAATGA GAATGGTGAG GGGAAATTGA 989  
 TTTCATTATT AAAACAAGG CTAGTTCTTC CTTTAGCATG GGAAGTGAG GTTGGGAGG 1049  
 GTAAGGACTA TAGCAGAATC TCTTCAATGA GCTTATTCCT TATCTTAGAC AAAACAGATT 1109  
 GTCACGGCAA GAGCAAGCAC TTGCTATATA ACCAAGTGCT TTCTCTTTTG CATTTTGAAAC 1169  
 AGCAATGGTC AGGCGCATG TGATTATGAA CTITTAACAC AGTAAACCCAG GTTCTTTTTC 1229  
 TGCCACATTT GCGAAGCTTC AGTGCAGCCT ATACCTTTTC ATAGCTGAG AAAATTAAGA 1289  
 GTATCCACTT ACTTAGATGG AAGAAATTAAT CAGTATAGAT TCTGATGACT CAGTTTGAAG 1349

CAGTGTCTCT CAACTGAAGC CCGCTGTGATA TTTTAAGAAA TATCTGGATT CCGAGGCTGG 1409  
 ACTCCTTTTT GTGGGCGAGT GTCCTGGGCA TTGTAGTAAT TTGGGAGCAC CCGTGGAGCT 1419  
 TAGCCACTAG ATACCAATAG CAGTCTCTCC CCAATGTAGT TGGGAGGAGC AGCCCAAAAT GTCTTCAGAC 1529  
 5 ACTGTCAAAAT GTGGCGAGGT GCGAAAATCA CCGCTGGTGG AGAACAGGGT CATCAATGCT 1589  
 AAGTACCTGT AACATTTTA ACTCTCAAAA CTGTGTATAT ACAAGGTCTA AATTATTAGA 1649  
 CGACCAATAC TTTAGTGTTA AAGGCATACA AATGAAATCT TCAAAAATCA AAATCTATTCT 1709  
 TGTTCCTCAA ATAGTGAATC TTATAAAAT AATCACAGAA GATGCAAAAT GATCAGAGT 1769  
 CCGTTAAAAAT TCGTCTCTGT ATGAGTATTT GAGGGAGGAA TTGGGTATAG TTCTCTACTT 1829  
 CTATTGGATG GTACTTTGAG ACTCAAAGC TAACTAAGT TGTGTGTGTG TCAGGCTGGC 1889  
 10 GGGTGTGGAA TCCACTCAGA TAAAGCAAA TCCATGTAAAT TGAATTCATA AGGTATATAT 1949  
 ATATCAAAAT GAAAAGTGGG CTATGACGCT TGGACAGCT AGAATTTTGA AAAATAATGG 2009  
 AATTCACAGG GATCTCTGAG AAGAGTACAC ATTTTACTCT GTGTACACTG CGACGACAGT 2129  
 CGACCCAGAA GACTCAGAAC AAAAGTACAC ATTTTACTCT GTGTACACTG CGACGACAGT 2129  
 GGGATTTAT TACTCTCTCC TCCCTAAAAA CCGACACAGC GGTCTCTCTT GGGAAATAAG 2189  
 AGGTTTCCAG CCGAAGAGA AGGAAGAGCT ATGTGTGTGT ACTCTAAAAA GTATTTAATA 2249  
 ACCGTTTTGT TGTGTGTGT GCTGTTTTGA AATCAGATTG TCCTCTCTCT ATATTTTATT 2309  
 15 TACTTCATTC TGTAAATTC TGTGGAATTA CTTAGAGCAA GCATGTGAA TTCTCAACTG 2369  
 TAAAGCAAAA TTCTCCATC ATTATAATTT CACATTTGTC CTGCGAGGT ATAAATTTTA 2429  
 TATTTCCACT GATAGTAATA AGGTAAATTC ATGACTCTAG TGGATAGATC TTTTCTATA 2489  
 AAGTACACT CATTATAGA GGGAGTCAAT GTTCACTGTC AGGAAGGTCA TTAGATAAAG 2549  
 CTCTGAATA TATTATGAAA CATTAGTCT GTTACTCTTA GATCTTTTT GTTAAATAAC 2609  
 TTTAAAGCT AACTTACCTA AAGAAATAT CTGACACATA TGAATCTCTC ATTAGAGTGC 2669  
 20 AGGAGAGAC CCAAGCCACA GATATGTAT TGAAGAAAGA ACAAGTCTC TAGGCCCGGC 2729  
 AGCTGGGCTC ACATCTGTAA TCTCAAGATG TTGAGAGGTC AAGGCCGGCA GATCACTGAA 2789  
 GCGTCAGAGT TCAAGACACG CCGTGGCAAC ATGATGAJAC CCGCTCTCTA CTAAATATC 2849  
 AAAAATATG AGGGATGTGT GGTGGAGGCG GTTGTGTGTA GCTGAGATCC CTCTACTGCA 2909  
 AGGAAATCT CTGATGAG CCAAGGGGAGG GTTGTGTGTA GCTGAGATCC CTCTACTGCA 2969  
 CTCGAGCTG GTGACAGAG ATGAGACTCG GTCCTCTGCG CCGCCCGCCG CTCTCCCCC 3029  
 25 AAAAAGATC TTCTTCATG AGAACATAG CCACTCAACA AAGGGAGACC TGGGTCTCAG 3089  
 TGTCCAAGTC ACTTATTCTG AGTAAATAG CAATGAAGA ATGCCATGGA ATCCCTGCCC 3149  
 AAATACCTCT GCTTATGATA TTGTAGAAAT TGATATAGAG TTGTATCCCA TTAAAGAGT 3209  
 AGGATGTAGT AGGAAATGAC TAAAAAACAAA CACACAAACA GAAACCCCT TTGTCTTGT 3269  
 AAGGTGGTTC CTAAGATAAT GTCACTGCAA TGTGGAAT AATATTAAAT ATGTGAAGT 3329  
 30 TTAGGCTTGT GTTTCCTCT CCGTCTCTCT TTCTCTGCCA GCGCTTGTCT ATTTTGCAG 3389  
 GTCAATGAAT CATGTAGAAA GAGACAGAG ATGAAGCTAG AACCACTAG AACCTGCCCT 3449  
 TTTTATATT TCTGTCTTGT GTAAAAGATA CAATGAGTGA GGAGGTGAG ATTTATAAT 3509  
 GAGTTTAAT AAGTTCTGT AGCTTTGATT TTCTCTTTC ATATTGTGA TCTTGCTATA 3569  
 GCCAGAAATG GCGCTGTAAAA TCTACATATG GATATTGAAG TCAAAATCTG TCAACTAGC 3629  
 35 TTACACTAGA TGGAGATATT TTCAATTTCA GATACACTGG AATGTATGAT CTAGCCATG 3689  
 GTAATATAGT CAAGTGTGTT AAGGTATTTA TTGTATAG CCGCTTGTGT TGGTACTG 3749  
 TTTCAAGTTT CTGCTCAATG ATTTCTTCAA ATTTTCAA TATTTTCCA TCAATGAGTA 3809  
 AAAATGCGCT GAGTCAACC TTCTGTGAAG TTGAACGAT CTGCTTTTTT AACAGTTTA 3869  
 AGCAATGGT ATATCATCTT CCGTTTACTA TGTAGCTTAA CTGAGGCTT ACCTTTTGA 3929  
 40 GTCAAGGCTC AACTTTATG CCACTTCAA AAGTTTATA TAATGTGTA AATTTTACT 3989  
 TCTCAAGGTT AGCATCTTA GAGTGTGCT CACAATTAG ATTCAGGAAA GAAAGACTT 4049  
 CAGTAGGAAC TGATTTGAAT TTAATGATG AGCATTCAT TGAATTTCTA GAATATAGT TATGAAGAT 4109  
 GATATTACAG CACAGACACA CGAGTATCT TGAATTTCTA GAATATAGT TATGAAGAT 4169  
 45 ATGGCTGACA CAGACGCTG ACTGCACTG AGCGAGGCT GAGTAAAGA ACAOCTCAT 4229  
 CTCTTCTCT TCTCTCTCAC ATTTCTAGAG CCGTTTGTAG GTAGCAAGA AATTGACTG 4289  
 CATTTGCAAT ACAAGGAGGA GAACTGTGCA AAGGGATGTA TGGTGGAGT TTTGTCTGT 4349  
 50 TTAATGAAGT GAAATATGAA AATGCTAGAG TTTTGTGCAA CATATAGTA GCAATAAAA 4409  
 CCAAGTGAAA AGTCTTTCCA AACTGTGTGT AAGAGGCGAT CTGCTGGGAA AAGATTGAG 4469  
 GAGAGGTAC TAAATTTGCT GGTATTTTCC GTAG G ACC CGA GAG AAT ACA 4523  
 Gly Thr Pro Glu Arg Asn Thr 115

VLT TGC AAA AGA TOT CCA GAT GGG TTC TTC TCA AAT GAG ACG TCA TCT 4571  
 Glt Cys Lys Arg Cys Pro Asp Gly Phe Phe Ser Asn Glu Thr Ser Ser 135

AAA GCA CCC TGT AGA AAA CAC ACA AAT TGC AGT GTC TTT GGT CTC CTG 4619  
 Lys Ala Pro Cys Arg Lys His Thr Asn Cys Ser Val Phe Gly Leu Leu 150

CTA ACT CAG AAA GGA AAT GCA ACA CAC GAC AAC ATA TGT TCC GGA AAC 4667



	Leu Thr Gln Lys Gly Asn Ala Thr His Asp Asn Ile Cys Ser Gly Asn	
	155 160 165	
5	AGT GAA TCA ACT CAA AAA TGT GGA ATA G GTAATTACAT TCCAAAATAC	4715
	Ser Glu Ser Thr Gln Lys Cys Gly Ile	
	170 175	
10	GTCTTTGTAC GATTTTGTAG TATCATCTCT CTCTCTGAGT TGAACACAAG GCCTCCAGCC	4775
	ACATTCCTGG TCAAACTTAC ATTTTCCCTT TCTTGAATCT TAACCAGTCA AGGCTACTCT	4835
	CGATGCATTA CTCTGAAGC TACCACCTCAG AATCTCTCAA AAATCTCACT CTCTACAGAT	4895
	AACACCTCAA AGCTTGATTT TCTCTCTTCT CACATCTGAA TCAAAATCTT CCAATAGGCA	4955
	AAGGCAGTG TCAATTTTCC CACTGAGATG AAATTAGGAG AGTCCAACT GTAGAAATCA	5015
	CGTTGTGTGT TATTACTTTC ACGAATGTCT GTATTATTAA CTAAAGTATA TATTGGCAAC	5075
	TAGAAGCAAA AGTGATATAA ACATGATGAC AAATTAGGCC AGGCATGGTG GCTTACTCT	5135
	ATAATCCCAA CATTTTGGGG GGCCAAAGTA GGCAGATCAC TTGAGGTGAG GATTCCAAG	5195
15	CCAGGCTGAC CAACATGGTG AAACCTGTCT TCTACTAAAA ATACAAAAAT TACGTGGGCA	5255
	TGTGTACAGG CACTTCTAGT ACCAGCTACT CAGGGCTGAG CGAGGAGGAT GCTTGAAGC	5315
	CAGAGATGG AGGTTGCAST GAGCTGAGT TGTACACGT CACTCCAGTG TGGGCAACAG	5375
	AGCAGATTT CATACACAC ACACACACAC ACACATTAGA AATGTGTACT	5435
	TGGCTTTGTT ACCATGGTA TTAGTGATCT TATTGCATGG AACTTCCAAG CTACTCTGT	5495
	TGTGTTAAGC TCTTCATTGG GTACAGGTCA CTAGATTAA GTTCAGGTTA TTGGATGCA	5555
	TTCCAGGTTA GTGATGACAA TTCACTAGGC TAGTGTGTGT GTTCACCTGT TCACTCCAC	5615
20	CACTAGACTA ATCTCAGACC TTCACTCAA GACACATFAC ACTAAAGATG ATTTGCTTCT	5675
	TGTGTTTAA TCAAGCAATG GTATAACCA GCTTGACTCT CCGCAACAG TTTTCCGAC	5735
	TACAAAGAG TTATGAAAG AGAGAAATCT GAATGTATA ATATATAGA TTCTAAOCCA	5795
	GTTCACGAT TGTTCATATG TGTAAATG ACATATAGA AGCCATTTA GCTTTCCTCT	5855
	CTCTATCTAA AAAAATAAAA AAAAATAATG AGGAAGGGGT ATTAAAGGA GTGATCAAT	5915
	TTTAACTTC TCTTAATTA ATTCATTTTT AATTACTT TTTTTCATT ATTGTGCACT	5975
25	TACTATGGG TACTGTGTA TAGAGGCTTT AACATTATA AAACACTGT GAAAGTGTCT	6035
	TACAGTAAAT ATAGGTAGTA GAAAGGACAG ACTAGATTTC AAAGCCAGGT CTAAGTAATC	6095
	CAAAAACAAA CACCCATTAC TCCCATTTTC TGGGAGATAC TTACTTACC CAGATGCTCT	6155
	GGGCTTGTA ATGCTATCTT AAATACATA GTTTATGTT TGTATATTT CTAATGTAAT	6215
	GTCTACTTAT ATATCTGAT CTATCTCTGT CTTTGTCTTC AAAGGTAAAC TATGTGTAA	6275
	AATGTGGGCA AAAAATAACA CACTATTCCA AATTACTGTT CAATTCCTCT TAAGTCAGTG	6335
	ATPATZATT GTTTTGACAT TAATCATGAA GTTCCCTGTG GGTACTAGGT AAACCTTTAA	6395
30	TAGATGTTA ATGTTTGTAT TCATTATAG AATTTTGGC GTTACTATAT TTACAACAAT	6455
	ATTTCACTCT AATTAGACAT TTACTAAACT TTCTCTGAA ACAATATGCC AAAAAGAAC	6515
	ATTAGAGAC ACGTAAAGTC AGTTGATCTT TGGCATAG ACAGAGCAC AGAAGCTTGA	6575
	TTTATTCGA ACTTGCAAT TTACGATAT TATCTTGA AAATCTCAAT GTGTGTGTT	6635
	TTTGTTTTG TTGTAATGA ATAGACTCTC AGAAATCCAA TTGTGAGTA AATCTTCTGG	6695
	GTTTCTAAC CTTCCTTTAG AT GTT ACC CTG TGT GAG GAG GCA TTC TTC ARG	6747
35	Asp Val Thr Leu Cys Glu Glu Ala Phe Phe Arg	
	180 185	
	TTT GCT GTT CCT ACA AAG TTT ACG CCT AAC TGG CTT AGT GTC TTG GTA	6795
	Phe Ala Val Pro Thr Lys Phe Thr Pro Asn Trp Leu Ser Val Leu Val	
	190 195 200	
40	GAC AAT TTG CCT GGC ACC AAA GTA AAC GCA GAG AGT GTA GAG AGG ATA	6843
	Asp Asn Leu Pro Gly Thr Lys Val Asn Ala Glu Ser Val Glu Arg Ile	
	205 210 215	
	AAA CGG CAA CAC AGC TCA CAA GAA CAG ACT TTC CAG CTG CTG AAG TTA	6891
	Lys Arg Gln His Ser Ser Gln Glu Gln Thr Phe Gln Leu Leu Lys Leu	
	220 225 230 235	
45	TGG AAA CAT CAA AAC AAA GAC CAA GAT ATA GTC AAG AAG ATC ATC CAA G	6940
	Trp Lys His Gln Asn Lys Asp Gln Asp Ile Val Lys Lys Ile Ile Gln	
	240 245 250	
50	GTATGATAAT CTAAATAAAA AAGATCAATC AGAAATCAA GACACCTATT TATCATAAAC	7000
	CAGGACAAG ACTCGATGTA TTTTAAAGTG TGTGAGTCT GTTCCCTGT TGAATCAATT	7060
	GTGGGACGTA AAAAGTTTCC ACTCGATAAT GTAGATGTA TTCCACAAC AGTTATACAA	7120
	GGTTTTGTTC TCACCCCTGC TCCCGAGTTT CTTGTAAAG TATGTTGAAC ACTCTAAGG	7180
	AAGGAAGATG CATTGAAGG CAGGGCTGTA TCTCAGGAG TCGCTCCAG ATCCCTTAAC	7240

GCTCTGTAA GCAGGCCCTC TAGACCACCA AGGAGAAGCT CTATAACCAC TTTGTATCTT 7300  
 ACATPTGCAOC TCTACCAAGA AGCTCTGTGT TATTTACTTG GTAATTTCTCT CCAGGTAGGC 7360  
 TTTTGTGAGC TTACAAATAT GTTCTTATTA ATCTCTCAAG TATGGGCTCG ATTAJAAATTA 7420  
 TTTTAATGGC ATATGTTATG AGAATTAATG AGATAAAATC TGAAAAGTGT TTGAGGCTCT 7480  
 TGTAGAGAAA AGCTAGTTAC AGCAAAATGT TCTACATCT TATAAGTTTA TATAAAGAT 7540  
 CTCCCTTAGA AATGGTGTGA GAGAGAAAACA GAGAGAGTA GCGAGAGAG TGTGAAGAA 7600  
 TCTGAADAAA AGAGATTCTA TCGATGTGTG ACTGTAAAGT TTACACACA TGTGTGAAG 7660  
 AGTCTGACT TCACTAAGCA TTGGGAGGAG ATGCTAGAG AAAAAAGAG AAGAGTTTCC 7720  
 ATAATGCGA CAGGCTCAGT GAGAAATTTA TTCAGGTCT CACCAGTAGT TAAATGACTG 7780  
 TATAGTCTTG CACTACCCTA AAAAAGTTCA AGTATCTGAA ACCGGGGCAA CAGATTTAG 7840  
 GAGACCAAG TCTTTGAGAG CTGATTGCTT TGCTTATGCA AAGAGATYAA CTTTATATGT 7900  
 TTGAGCAAG CAAAATATTT CTTTGAACCT ATAAATAGCC CTGAGGCCGA AGAGAAAGG 7960  
 AAAAAAGAG ACCGTTAGAA TTGGAAGCAA CCAATTCCT TATTTTTAA ATGAGGACAT 8020  
 TTAAAGCCAG AAAAGTGAAC CGATTTGGCT TAGGGCTCAC AGATACTAG TGACTCATGT 8080  
 CATTAATAGA AATGTTAGTT CTTCCCTCTT AGGTTTGTAC CTAAGCTTAT TACTGAATA 8140  
 TTCTTAGGC TGTGTGCTC CTTTAGTTCC TGCACTCAT GTCTTTGAGT TTTGAGTAT 8200  
 CTTCTCATG GAGGTAGTCC TCTGGTGCTA TGTGTATCT TAAAGGCTA GTTACGGCAA 8260  
 TTAATTTAT AACTAGCGCC TACTAATGAA ACTTGTATTT ACAGAGTAGT TACTTTGAT 8320  
 ACTTTCTTT TTTTCTGAAA TGTATGGTG GTAATTCTCT AACTTTTCT TGAJAAACT 8380  
 GAGAGTTAG TGTCTATTT TCTACGTTTA ATTTCAAAA TATGAGGCTT CTTCCAAGT 8440  
 TTGTGTGAGT GCGAAJATA TATGACATAT TATCTATTA TACCAAAAA TATTTATCTC 8500  
 AGTCTTAGA AATAAATGCT GTCACTTAAC TCGCTCTCAA AAGAAAAGGT TATCATTTGA 8560  
 ATATAATTAT GAATTTCTCG AAGAACCTTT TGCTCAAGC TTGTTTTATG ATGGCATTTG 8620  
 ATGAATATA ATGATGTGAA CACTTATCTG GCTTTTGTCT TATGCAAT AT ATT GAC 8675  
 Asp Ile Asp

CTC TGT GAA AAC AGC GTG CAG CGG CAC ATT GGA CAT GCT AAC CTC ACC 8724  
 Leu Cys Glu Asn Ser Val Gln Arg His Ile Gly His Ala Asn Leu Thr  
 255 260 265 270

TTC GAG CAG CTT CGT AGC TTG ATG GAA AGC TTA CCG GGA AAG AAA GTG 8772  
 Phe Glu Gln Leu Arg Ser Leu Met Glu Ser Leu Pro Gly Lys Lys Val  
 275 280 285

GGA GCA GAA GAC ATT GAA AAA ACA ATA AAG GCA TGC AAA OCC AGT GAC 8820  
 Gly Ala Glu Asp Ile Glu Lys Thr Ile Lys Ala Cys Lys Pro Ser Asp  
 290 295 300

CAG ATC CTG AAG CTG CTC AGT TTG TGG CGA ATA AAA AAT GGC GAC CAA 8868  
 Gln Ile Leu Lys Leu Ser Leu Trp Arg Ile Lys Asn Gly Asp Gln  
 305 310 315

GAC ACC TTG AAG GGC CTA ATG CAC GCA CTA AAG CAC TCA AAG AGG TAC 8916  
 Asp Thr Leu Lys Gly Leu Met His Ala Leu Lys His Ser Lys Thr Tyr  
 320 325 330

CAC TTT OCC AAA ACT GTC ACT CAG AGT CTA AAG AAG ACC ATC AGG TTC 8964  
 His Phe Pro Lys Thr Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe  
 335 340 345 350

CTT CAC AGC TTC ACA ATG TAC AAA TTG TAT CAG AAG TTA TTT TTA GAA 9012  
 Leu His Ser Phe Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu  
 355 360 365

ATG ATA GGT AAC CAG GTC CAA TCA GTA AAA ATA AGC TGC TTA 9054  
 Met Ile Gly Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu  
 370 375 380

TAACTGGAAA TGCCATTGA GCTGTTTCT CACAATTGGC GAGATCCAT GGTAGTAAA 9114  
 ACTGTTTCTC AGGCACCTTGA GGCCTTTAGT GATATCTTC TCATTACAG TCATATATT 9174  
 TGCCACAGGG TACTAAAAAG AACATGATG TGGAGAAAG ACTACATCT CTCCAAATA 9234  
 ACCCAAAAT GTTAAATCAA CTGTACAGAT TGATCGTTA TCTACTGAT ATTTTTCCT 9294  
 TTTACTGTC TTGCAGTAAT TCAACTGGAA ATTAAAAAA AAAAATGAT CTCCACTGGG 9354  
 CTTTACTAAA TATGGGAATG TCAACTTAA ATAGCTTGG GATTCCAGCT ATGCTAGAGG 9414  
 CTTTATTAG AAAGCCATAT TTTTTCCTG AAAAGTACT AATATCTG TAACATATT 9474

ACAGTATTGC TATTATATT CATTAGATA TAAGATTGG ACATATTATC ATCCTATAAA 9514  
 GAAACGGTAT GACTTAATTT TAGAAGAAA ATTATATCTT GTTATATATG ACAAATGAAA 9594  
 GAGAAATAT ATATTTTAA TGGAAAGTTT GTAGCATTTT TCTAATAGTG ACTGCCATAT 9654  
 TTTCGCTGT GGAGTATTTT TATAATTTA TCTGTATAG CTGTATATC ATTTTATAGA 9714  
 AAATGATA TTATGCAAT TGTTAATGT TGGAAACT ATGAAATATA AATATCTGA 9774  
 ATATTAGATG CACTGAGAAA TTGAATGTAC CTATTTAAA AGATTATTATG GTTTTATAAC 9814  
 TATATAATG ACATTATTA AGTTTTCAAA TTATTTTAA TTGCTTCTC TGTGCTTTT 9898  
 ATTT

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 401 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Asn Asn Leu Leu Cys Cys Ala Leu Val Phe Leu Asp Ile Ser  
 -20 -15 -10  
 Ile Lys Trp Thr Thr Gln Glu Thr Phe Pro Pro Lys Tyr Leu His  
 5 1 5  
 Tyr Asp Glu Glu Thr Ser His Gln Leu Leu Cys 5 Asp Lys Cys Pro  
 10 15 20  
 Pro Gly Thr Tyr Thr Leu Lys Gln His Cys Thr Ala Lys Trp Lys Thr  
 25 30 35  
 Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp Ser Trp His  
 40 45 50  
 Thr Ser Asp Glu Cys Leu Tyr Cys Ser Pro Val Cys Lys Glu Leu  
 55 60 65  
 Gln Tyr Val Lys Gln Glu Cys Asn Arg Thr His Asn Arg Val Cys  
 70 75 80  
 Glu Cys Lys Glu Gly Arg Tyr Leu Glu Ile Glu Phe Cys Leu Lys  
 85 90 95  
 His Arg Ser Cys Pro Pro Gly Phe Gly Val Val Gln Ala Gly Thr  
 100 105 110  
 Pro Glu Arg Asn Thr Val Cys Lys Arg Cys Pro Asp Gly Phe Phe  
 115 120 125  
 Ser Asn Glu Thr Ser Ser Lys Ala Pro Cys Arg Lys His Thr Asn  
 130 135 140  
 Cys Ser Val Phe Gly Leu Leu Leu Thr Gln Lys Gly Asn Ala Thr  
 145 150 155  
 His Asp Asn Ile Cys Ser Gly Asn Ser Glu Ser Thr Gln Lys Cys  
 160 165 170  
 Gly Ile Asp Val Thr Leu Cys Glu Glu Ala Phe Phe Arg Phe Ala  
 175 180 185  
 Val Pro Thr Lys Phe Thr Pro Asn Trp Leu Ser Val Leu Val Asp  
 190 195 200  
 Asn Leu Pro Gly Thr Lys Val Asn Ala Glu Ser Val Glu Arg Ile  
 205 210 215  
 Lys Arg Gln His Ser Ser Gln Glu Gln Thr Phe Gln Leu Leu Lys  
 220 225 230  
 Leu Trp Lys His Gln Asn Lys Asp Gln Asp Ile Val Lys Lys Ile  
 235 240 245  
 Ile Gln Asp Ile Asp Leu Cys Glu Asn Ser Val Gln Arg His Ile  
 250 255 260  
 Gly His Ala Asn Leu Thr Phe Glu Gln Leu Arg Ser Leu Met Glu  
 265 270 275  
 Ser Leu Pro Gly Lys Lys Val Gly Ala Glu Asp Ile Glu Lys Thr  
 280 285 290  
 Ile Lys Ala Cys Lys Pro Ser Asp Gln Ile Leu Lys Leu Leu Ser  
 295 300 305

Leu Trp Arg Ile Lys Asn Gly Asp Gln Asp Thr Leu Lys Gly Leu  
 310 315 320  
 Met His Ala Leu Lys His Ser Lys Thr Tyr His Phe Pro Lys Thr  
 325 330 335  
 Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe Leu His Ser Phe  
 340 345 350  
 Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu Met Ile Gly  
 355 360 365  
 Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu  
 370 375 380

## (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1206 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ATGAACAAC TGCCTGCTG CGCGCTCGTG TTTCTGGACA TCTCCATTAA GTGGACCACC 60  
 CAGGAACAGT GTCTCCCAA GTACCTTCAT TATGACGAAG AACCTCTCA TCAGCTGTGT 120  
 TGTGACAAAT GTCTCCCTGG TACCTACCTA AAACAACACT GTACAGCAAA GTGGAAAGACC 180  
 GTGTGCGCCC CTTGCCCTGA CCACTACTAC ACAGACAGCT GGCACACCAG TGACGAGTGT 240  
 CTATACTGCA GCCCGGTGTG CAAGGAGCTG CAGTACGTCA AGCAGGAGTG CAATCGCACC 300  
 CACACCGCGG TGTGCGAATG CAAGGAAGGG CGCTACCTTG AGATAGAGTT CTGCTTGAAA 360  
 CATAGGAGCT GCCCTCCCTG ATTGGAGTG GTGCAAGCTG GAACCCAGAG GCGAAATACA 420  
 GTTTGCAAAA GATGTCACGA TGGGTCTCTC TCAAAATGAG CGTCATCTAA AGCACCCCTGT 480  
 AGAAAACACA CAAATTCGAG TGTCTTTGGT CTCCTGCTAA CTCAGAAAGG AAATGCACAA 540  
 CACGACAACA CATGTTCCGG AAACAGTGAA TCAACTCAAA AATGTGGAAT AGATGTTACC 600  
 CTGTGTGAGG AGGCATCTCT CAGGTTTGCT GTTCTCAAAA AGTTTACGCC TAACCTGGCTT 660  
 AGTGTCTTGG TAGACAATTT GCCTGGCACC AAAGTAAACG CAGAGAGTGT AGAGAGGATA 720  
 AAACGCAAC ACAGCTCACA AGAACAGACT TTCCAGCTGC TGAAGTTATG GAAACATCAA 780  
 AGTGTCTTGG TAGACAATTT GCCTGGCACC AAAGTAAACG CAGAGAGTGT AGAGAGGATA 840  
 GTGCAAGGCG ACATTGGACA TGCTAACCTC ACCTTCGAGC AGCTTCTGAG CTTGATGGAA 900  
 AGCTTACCGG GAAAGAAAGT GGGAGCAGAA GACATTGAAA AAACAATAAA GGCAATGCAA 960  
 CCAAGTAGAC AGATCTCTGAA GCTGCTCAGT TTTGTGGCAA TAAAAATATG CGACCAAGAC 1020  
 ACCTTGAAGG GCCTTAATGCA CGCACTAAGG CACTCAAAAG CTGACCACTT TCCCAAACT 1080  
 GTCACTCAGA GTCTAAAGAA GACCATCAGG TTTCTTCACA GCTTCACAA GTACAAATG 1140  
 TATCAGAAAG TATTTTGTGA AATGATAGGT AACCAAGTCC AATCAGTAAA AATAAGCTGC 1200  
 TTATAA 1206

## Claims

1. A DNA comprising the nucleotide sequences of the Sequences No. 1 and No. 2 in the Sequence Table.
2. The DNA according to claim 1, wherein the Sequence ID No. 1 includes the first exon of the OCIF gene and the Sequence ID No. 2 includes the second, third, fourth, and fifth exons.
3. A protein exhibiting the activity of inhibiting differentiation and/or maturation of osteoclasts and having the following physicochemical characteristics,

(a) molecular weight (SDS-PAGE):

- (i) Under reducing conditions: about 60 kD,
- (ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 in the Sequence Table,

(c) affinity:

exhibits affinity to a cation exchanger and heparin, and

(d) heat stability:

- (i) the osteoclastogenesis-inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,
- (ii) the osteoclastogenesis-inhibitory activity is lost when treated with heat at 90°C for 10 minutes.

4. A process for producing a protein exhibiting an activity of inhibiting differentiation and/or maturation of osteoclasts and having the following physicochemical characteristics,

(a) molecular weight (SDS-PAGE):

- (i) Under reducing conditions: about 60 kD,
- (ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 of the Sequence Table,

(c) affinity:


exhibits affinity to a cation exchanger and heparin, and

(d) heat stability:

- (i) the osteoclastogenesis-inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,
- (ii) the osteoclastogenesis-inhibitory activity is lost when treated with heat at 90°C for 10 minutes,

the process comprising inserting a DNA including the nucleotide sequences of the sequences No. 1 and No. 2 in the Sequence Table into an expression vector, producing a vector capable of expressing a protein having the above-mentioned physicochemical characteristics and exhibiting the activity of inhibiting differentiation and/or maturation of osteoclasts, and producing this protein by a genetic engineering technique.

Figure 1



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP97/02859

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int. C1 <sup>6</sup> C12N15/00, C12P21/00 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) Int. C1 <sup>6</sup> C12N15/00, C12P21/00 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WPI, GENETIX-CDROM, BIOSIS		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Cancer Research, (1995), Vol. 55, Toshiyuki Yoneda, et al. "Sumarin suppresses hypercalcemia and osteoclastic bone resorption in nude mice bearing a human squamous cancer" P. 1989-1993	1 - 4
A	Proc. Natl. Acad. Sci. USA, (1990) Vol. 87 Kukita A. et al. "Osteoinductive factor inhibits formation of human osteoclast-like cells" P. 3023-3026	1 - 4
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date: document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another claim or other special reason (as specified) "O" document referring to oral disclosure, use, exhibition or other means "T" document published prior to the international filing date but later than the priority date claimed "X" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "Y" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Z" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family		
Date of the actual completion of the international search September 29, 1997 (29. 09. 97)		Date of mailing of the international search report October 7, 1997 (07. 10. 97)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA210 (second sheet) (July 1992)